

## TECHNICAL DATA SHEET

### CyFlow™ CD107b APC Anti-Hu; Clone H4B4

**REF** CT778572

**For Research Use Only.**  
**Not for use in diagnostic or therapeutic procedures.**

### Specifications

|                                    |                                                 |
|------------------------------------|-------------------------------------------------|
| <b>Antigen</b>                     | CD107b                                          |
| <b>Alternative Names</b>           | LAMP-2                                          |
| <b>Clone</b>                       | H4B4                                            |
| <b>Clonality</b>                   | monoclonal                                      |
| <b>Format</b>                      | APC                                             |
| <b>Host / Isotype</b>              | Mouse / IgG1                                    |
| <b>Species Reactivity</b>          | Human                                           |
| <b>Negative Species Reactivity</b> | Mouse   Rat                                     |
| <b>Quantity</b>                    | 100 tests                                       |
| <b>Immunogen</b>                   | Human peripheral blood mononuclear cells (PBMC) |

### Specificity

The mouse monoclonal antibody H4B4 recognizes CD107b antigen, an extensively glycosylated 100-120 kDa widely expressed lysosome-associated protein.

#### Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)

## Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

## Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.02% (w/v) sodium azide.

## Storage and Stability

|                  |                                                                               |
|------------------|-------------------------------------------------------------------------------|
| <b>Storage</b>   | Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze. |
| <b>Stability</b> | Do not use after expiration date stamped on vial label.                       |

## Background Information

CD107b (LAMP-2; lysosome-associated membrane protein-2), together with CD107a (LAMP-1), is a major constituent of lysosomal membrane. The LAMP proteins are involved in lysosome biogenesis and are required for fusion of lysosomes with phagosomes, especially CD107b is important regulator in successful phagosomal maturation. CD107b deficiency causes an accumulation of autophagosomes in many tissues leading to cardiomyopathy and myopathy (Danons disease). Immature CD107b is an approximately 45 kDa protein, but after extensive glycosylation the mature glycoprotein has about 100-120 kDa.

## References

- Denzer K, van Eijk M, Kleijmeer MJ, Jakobson E, de Groot C, Geuze HJ: Follicular dendritic cells carry MHC class II-expressing microvesicles at their surface. J Immunol. 2000 Aug 1; 165(3):1259-65. < PMID: 10903724 >
- Palmer DR, Fernandez S, Bisbing J, Peachman KK, Rao M, Barvir D, Gunther V, Burgess T, Kohno Y, Padmanabhan R, Sun W: Restricted replication and lysosomal trafficking of yellow fever 17D vaccine virus in human dendritic cells. J Gen Virol. 2007 Jan; 88(1):148-56. < PMID: 17170447 >
- Borgne-Sanchez A, Dupont S, Langonné A, Baux L, Lecoeur H, Chauvier D, Lassalle M, Déas O, Brière JJ, Brabant M, Roux P, Péchoux C, Briand JP, Hoebeke J, Deniaud A, Brenner C, Rustin P, Edelman L, Rebouillat D, Jacotot E: Targeted Vpr-derived peptides reach mitochondria to induce

---

### Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)

- apoptosis of alphaVbeta3-expressing endothelial cells. Cell Death Differ. 2007 Mar; 14(3):422-35. < PMID: 16888644 >
- Kannanganat S, Ibegbu C, Chennareddi L, Robinson HL, Amara RR: Multiple-cytokine-producing antiviral CD4 T cells are functionally superior to single-cytokine-producing cells. J Virol. 2007 Aug; 81(16):8468-76. < PMID: 17553885 >
  - Guia S, Cognet C, de Beaucoudrey L, Tessmer MS, Jouanguy E, Berger C, Filipe-Santos O, Feinberg J, Camcioglu Y, Levy J, Al Jumaah S, Al-Hajjar S, Stephan JL, Fieschi C, Abel L, Brossay L, Casanova JL, Vivier E: A role for interleukin-12/23 in the maturation of human natural killer and CD56+ T cells in vivo. Blood. 2008 May 15; 111(10):5008-16. < PMID: 18319400 >
  - Apte SH, Baz A, Groves P, Kelso A, Kienzle N: Interferon-gamma and interleukin-4 reciprocally regulate CD8 expression in CD8+ T cells. Proc Natl Acad Sci USA. 2008 Nov 11; 105(45):17475-80. < PMID: 18988742 >
  - Thedrez A, Harly C, Morice A, Salot S, Bonneville M, Scotet E: IL-21-mediated potentiation of antitumor cytolytic and proinflammatory responses of human V gamma 9V delta 2 T cells for adoptive immunotherapy. J Immunol. 2009 Mar 15; 182(6):3423-31. < PMID: 19265120 >
  - Meade JL, Wilson EB, Holmes TD, de Wynter EA, Brett P, Straszynski L, Ballard PA, Trapani JA, McDermott MF, Cook GP: Proteolytic activation of the cytotoxic phenotype during human NK cell development. J Immunol. 2009 Jul 15; 183(2):803-13. < PMID: 19570824 >

---

The Safety Data Sheet for this product is available at [www.sysmex-partec.com/services](http://www.sysmex-partec.com/services).

---

---

**Contact Information:**

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany  
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: [info@sysmex-partec.com](mailto:info@sysmex-partec.com)