

TECHNICAL DATA SHEET

CyFlow™ CD11b APC Anti-Hu/Ms; Clone M1/70

REF BR751578

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD11b
Alternative Names	Mac-1
Clone	M1/70
Clonality	monoclonal
Format	APC
Host / Isotype	Rat / IgG2b
Species Reactivity	Human Mouse, Non-Human Primates Rabbit
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [0.5 mg/ml]
Immunogen	B10 mouse spleen cells enriched for T cells

Specificity

The rat monoclonal antibody M1/70 recognizes CD11b antigen, a type I transmembrane protein mainly expressed on monocytes/macrophages, granulocytes and NK-cells, which associates with CD18 to form Mac-1 integrin that plays important role in cell-cell interactions.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Working concentrations should be determined by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD11b (integrin α M subunit) is a 165-170 kDa type I transmembrane glycoprotein that non-covalently associates with CD18 (integrin β 2 subunit); expression of the CD11b chain on the cell surface requires the presence of the CD18 antigen. CD11b/CD18 integrin (Mac-1, CR3) is highly expressed on NK cells, neutrophils, monocytes and less on macrophages. CD11b/CD18 integrin is implicated in various adhesive interactions of monocytes, macrophages and granulocytes, facilitating their diapedesis, as well as it mediates the uptake of complement coated particles, serving as a receptor for the iC3b fragment of the third complement component.

References

- Springer T, Galfrè G, Secher DS, Milstein C: Monoclonal xenogeneic antibodies to murine cell surface antigens: identification of novel leukocyte differentiation antigens. Eur J Immunol. 1978 Aug; 8(8):539-51. < PMID: 81133 >
- Ault KA, Springer TA: Cross-reaction of a rat-anti-mouse phagocyte-specific monoclonal antibody (anti-Mac-1) with human monocytes and natural killer cells. J Immunol. 1981 Jan; 126(1):359-64. < PMID: 7451976 >
- Sanchez-Madrid F, Simon P, Thompson S, Springer TA: Mapping of antigenic and functional epitopes on the alpha- and beta-subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. J Exp Med. 1983 Aug 1; 158(2):586-602. < PMID: 6193226 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Whiteland JL, Nicholls SM, Shimeld C, Easty DL, Williams NA, Hill TJ: Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. *J Histochem Cytochem.* 1995 Mar; 43(3):313-20. < PMID: 7868861 >
- Welt FG, Edelman ER, Simon DI, Rogers C: Neutrophil, not macrophage, infiltration precedes neointimal thickening in balloon-injured arteries. *Arterioscler Thromb Vasc Biol.* 2000 Dec; 20(12):2553-8. < PMID: 11116052 >
- Dembic Z, Schenck K, Bogen B: Dendritic cells purified from myeloma are primed with tumor-specific antigen (idiotype) and activate CD4+ T cells. *Proc Natl Acad Sci USA.* 2000 Mar 14; 97(6):2697-702. < PMID: 10706628 >
- Zhang Y, McCormick LL, Desai SR, Wu C, Gilliam AC: Murine sclerodermatous graft-versus-host disease, a model for human scleroderma: cutaneous cytokines, chemokines, and immune cell activation. *J Immunol.* 2002 Mar 15; 168(6):3088-98. < PMID: 11884483 >
- Brickson S, Ji LL, Schell K, Olabisi R, St Pierre Schneider B, Best TM: M1/70 attenuates blood-borne neutrophil oxidants, activation, and myofiber damage following stretch injury. *J Appl Physiol.* 2003 Sep; 95(3):969-76. < PMID: 12730143 >
- Leon F, Contractor N, Fuss I, Marth T, Lahey E, Iwaki S, la Sala A, Hoffmann V, Strober W, Kelsall BL: Antibodies to complement receptor 3 treat established inflammation in murine models of colitis and a novel model of psoriasiform dermatitis. *J Immunol.* 2006 Nov 15; 177(10):6974-82. < PMID: 17082612 >
- Takagi H, Numazaki M, Kajiwara T, Abe Y, Ishii M, Kato C, Kojima N: Cooperation of specific ICAM-3 grabbing nonintegrin-related 1 (SIGNR{01} and complement receptor type 3 (CR3) in the uptake of oligomannose-coated liposomes by macrophages. *Glycobiology.* 2009 Mar; 19(3):258-66. < PMID: 19029201 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com