

TECHNICAL DATA SHEET

CyFlow™ CD147 Purified Anti-Hu; Clone MEM-M6/2

REF AS261764

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD147
Alternative Names	Basigin, BSG, M6, EMMPRIN, 5F7, TCSF
Clone	MEM-M6/2
Clonality	monoclonal
Format	Purified
Host / Isotype	Mouse / IgG2b
Species Reactivity	Human, Pig
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	A soluble recombinant form of CD147, CD147Rg, which consists of the cDNA coding for the entire extracellular region of CD147 fused to the DNA coding for the hinge region, CH2 and CH3 domain of human IgG1

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Specificity

The mouse monoclonal antibody MEM-M6/2 recognizes extracellular Ig domain D1 of CD147 antigen, a 50-60 kDa type I transmembrane glycoprotein primarily expressed on all leukocytes, red blood cells, platelets and endothelial cells; it is not expressed by resting lymphocytes. The antibody MEM-M6/1 is a high-affinity antibody capable of binding to unstimulated peripheral blood T cells.

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunohistochemistry (paraffin-embedded sections)

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD147 (basigin, neurothelin, OX-47, 5A11, CE9, M6) also known as EMMPRIN (extracellular matrix metalloproteinase inducer) or TCSF (tumour cell-derived collagenase-stimulatory factor) is an ubiquitously expressed cell surface protein with multiple glycosylated forms. The highest level of CD147 expression is on metabolically active cells, such as lymphoblasts, inflammatory cells, brown adipocytes and malignant tumor cells. CD147 has multiple functions, including facilitating of cell surface expression of monocarboxylate transporter proteins and extracellular matrix metalloproteinases, regulation of integrin functions, it plays roles in cell development and activation, fetal development or retinal function.

References

- Koch C, Staffler G, Huttinger R, Hilgert I, Prager E, Cerny J, Steinlein P, Majdic O, Horejsi V, Stockinger H: T cell activation-associated epitopes of CD147 in regulation of the T cell response, and their definition by antibody affinity and antigen density. Int Immunol. 1999 May; 11(5):777-86. < PMID: 10330283 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Kirk P, Wilson MC, Heddle C, Brown MH, Barclay AN, Halestrap AP: CD147 is tightly associated with lactate transporters MCT1 and MCT4 and facilitates their cell surface expression. EMBO J. 2000 Aug 1; 19(15):3896-904. < PMID: 10921872 >
- Wilson MC, Meredith D, Fox JE, Manoharan C, Davies AJ, Halestrap AP: Basigin (CD147 is the target for organomercurial inhibition of monocarboxylate transporter isoforms 1 and 4: the ancillary protein for the insensitive MCT2 is EMBIGIN (gp70). J Biol Chem. 2005 Jul 22; 280(29):27213-21. < PMID: 15917240 >
- Xu D, Hemler ME: Metabolic activation-related CD147-CD98 complex. Mol Cell Proteomics. 2005 Aug; 4(8):1061-71. < PMID: 15901826 >
- Iacono KT, Brown AL, Greene MI, Saouaf SJ: CD147 immunoglobulin superfamily receptor function and role in pathology. Exp Mol Pathol. 2007 Dec; 83(3):283-95. < PMID: 17945211 >
- Melchior A, Denys A, Deligny A, Mazurier J, Allain F: Cyclophilin B induces integrin-mediated cell adhesion by a mechanism involving CD98-dependent activation of protein kinase C-delta and p44/42 mitogen-activated protein kinases. Exp Cell Res. 2008 Feb 1; 314(3):616-28. < PMID: 18054915 >
- Schmidt R, Bültmann A, Fischel S, Gillitzer A, Cullen P, Walch A, Jost P, Ungerer M, Tolley ND, Lindemann S, Gawaz M, Schömig A, May AE: Extracellular matrix metalloproteinase inducer (CD147 is a novel receptor on platelets, activates platelets, and augments nuclear factor kappaB-dependent inflammation in monocytes. Circ Res. 2008 Feb 15; 102(3):302-9. < PMID: 18048771 >
- Ruiz S, Castro-Castro A, Bustelo XR: CD147 Inhibits the Nuclear Factor of Activated T-cells by Impairing Vav1 and Rac1 Downstream Signaling. J Biol Chem. 2008 Feb 29; 283(9):5554-66. < PMID: 18160397 >
- Plánka L, Necas A, Srnc R, Rauser P, Starý D, Jancár J, Amler E, Filová E, Hlucilová J, Kren L, Gál P: Use of allogenic stem cells for the prevention of bone bridge formation in miniature pigs. Physiol Res. 2009; 58(6):885-93. < PMID: 19093735 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com