

TECHNICAL DATA SHEET

CyFlow™ CD16 Purified Anti-Hu; Clone 3G8

REF BM323481

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD16
Alternative Names	FcγRIIIA, CD16a
Clone	3G8
Clonality	monoclonal
Format	Purified
Host / Isotype	Mouse / IgG1
Species Reactivity	Human, Non-Human Primates
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Human neutrophils

Specificity

The mouse monoclonal antibody 3G8 recognizes CD16 antigen, a low affinity receptor for aggregated IgG (FcγRIII antigen). CD16 exists in two different isoforms: CD16a (FcγRIIIA; 50-65 kDa; expressed on

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

NK-cells, monocytes and macrophages) and CD16b (FcγRIIIB; 48 kDa; mainly expressed on neutrophils).

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Immunohistochemistry (frozen sections)
- Functional assays

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD16 (FcγRIII) is a 50-65 kDa glycoprotein serving as a low affinity IgG receptor. Human FcγRIII is expressed in two forms - FcγRIIIA and FcγRIIIB. FcγRIIIA is a transmembrane protein of monocytes, macrophages, NK cells and a subset of T cells. It is associated with FcεRI-γ subunit and is responsible for antibody-dependent NK cell cytotoxicity. Mast cell FcγRIIIA is associated, moreover, with FcεRI-β subunit. Besides IgG, FcγRIIIA can be triggered also by oligomeric IgE. FcγRIIIB is a GPI-linked monomeric receptor expressed on neutrophils and is involved in their activation and induction of a proadhesive phenotype.

References

- Knapp W, Dorken B, Gilks W, Rieber EP, Schmidt RE, Stein H, von dem Borne AEGK (Eds): Leucocyte Typing IV. Oxford University Press, Oxford. 1989; 1-1820. < NLM ID: 8914679 >
- Doussis IA, Gatter KC, Mason DY: CD68 reactivity of non-macrophage derived tumours in cytological specimens. J Clin Pathol. 1993 Apr; 46(4):334-6. < PMID: 7684403 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Metes D, Ernst LK, Chambers WH, Sulica A, Herberman RB, Morel PA: Expression of functional CD32 molecules on human NK cells is determined by an allelic polymorphism of the FcγRIIc gene. *Blood*. 1998 Apr 1; 91(7):2369-80. < PMID: 9516136 >
- Zhu X, Hamann KJ, Muñoz NM, Rubio N, Mayer D, Herrnreiter A, Leff AR: Intracellular expression of Fc γRIII (CD16) and its mobilization by chemoattractants in human eosinophils. *J Immunol*. 1998 Sep 1; 161(5):2574-9. < PMID: 9725258 >
- Wijngaarden S, van Roon JA, van de Winkel JG, Bijlsma JW, Lafeber FP: Down-regulation of activating Fcγ receptors on monocytes of patients with rheumatoid arthritis upon methotrexate treatment. *Rheumatology (Oxford)*. 2005 Jun; 44(6):729-34. < PMID: 15757966 >
- Komano Y, Nanki T, Hayashida K, Taniguchi K, Miyasaka N: Identification of a human peripheral blood monocyte subset that differentiates into osteoclasts. *Arthritis Res Ther*. 2006; 8(5):R152. < PMID: 16987426 >
- Congy-Jolivet N, Bolzec A, Ternant D, Ohresser M, Watier H, Thibault G: Fc γRIIIa expression is not increased on natural killer cells expressing the Fc γRIIIa-158V allotype. *Cancer Res*. 2008 Feb 15; 68(4):976-80. < PMID: 18281470 >
- Choi EI, Wang R, Peterson L, Letvin NL, Reimann KA: Use of an anti-CD16 antibody for in vivo depletion of natural killer cells in rhesus macaques. *Immunology*. 2008 Jun; 124(2):215-22. < PMID: 18201184 >
- Burt BM, Plitas G, Zhao Z, Bamboat ZM, Nguyen HM, Dupont B, DeMatteo RP: The lytic potential of human liver NK cells is restricted by their limited expression of inhibitory killer Ig-like receptors. *J Immunol*. 2009 Aug 1; 183(3):1789-96. < PMID: 19587011 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com