

TECHNICAL DATA SHEET

CyFlow™ CD178 Low Endotoxin Anti-Hu; Clone NOK-1

REF BG866777

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD178
Alternative Names	Fas-L, FAS-Ligand, CD95L, CD95-L, APO-1L, APT1LG, TNFSF6 ligand
Clone	NOK-1
Clonality	monoclonal
Format	Low Endotoxin
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	L5178Y mouse T lymphoma cells expressing recombinant human CD178

Specificity

The mouse monoclonal antibody NOK-1 recognizes CD178 antigen, a 40 kDa transmembrane glycoprotein expressed on neutrophils, monocytes, and activated T and NK cells.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Western blot
- Immunocytochemistry
- Functional assays

Storage Buffer

The reagent is provided in azide-free phosphate buffered saline (PBS) solution, pH ≈7.4; 0.2 µm filter sterilized. Endotoxin level is less than 0.01 EU/µg of the protein, as determined by the LAL test.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD178 (Fas-L; Fas ligand or CD95L), a member of TNF family transmembrane glycoproteins, is responsible for induction of apoptosis in cells containing its receptor Fas/CD95. The CD178-mediated apoptosis pathway has been implicated in peripheral tolerance, tissue pathology, and maintenance of the immune privileged sites. Defects in this interaction may be related to some cases of systemic lupus erythematosus (SLE). CD178 was also described as a co-stimulatory receptor for T-cell activation in mice in vivo.

References

- Kayagaki N, Kawasaki A, Ebata T, Ohmoto H, Ikeda S, Inoue S, Yoshino K, Okumura K, Yagita H: Metalloproteinase-mediated release of human Fas ligand. J Exp Med. 1995 Dec 1; 182(6):1777-83. < PMID: 7500022 >
- Ghadimi MP, Sanzenbacher R, Thiede B, Wenzel J, Jing Q, Plomann M, Borkhardt A, Kabelitz D, Janssen O: Identification of interaction partners of the cytosolic polyproline region of CD95 ligand (CD178). FEBS Lett. 2002 May 22; 519(1-3):50-8. < PMID: 12023017 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Chopin D, Barei-Moniri R, Maillé P, Le Frère-Belda MA, Muscatelli-Groux B, Merendino N, Lecerf L, Stoppacciaro A, Velotti F: Human urinary bladder transitional cell carcinomas acquire the functional Fas ligand during tumor progression. Am J Pathol. 2003 Apr; 162(4):1139-49. < PMID: 12651606 >
- Jodo S, Pidiyar VJ, Xiao S, Furusaki A, Sharma R, Koike T, Ju ST: Fas ligand (CD178) cytoplasmic tail is a positive regulator of Fas ligand-mediated cytotoxicity. J Immunol. 2005 Apr 15; 174(8):4470-4. < PMID: 15814666 >
- Ji J, Chen JJ, Braciale VL, Cloyd MW: Apoptosis induced in HIV-1-exposed, resting CD4+ T cells subsequent to signaling through homing receptors is Fas/Fas ligand-mediated. J Leukoc Biol. 2007 Jan; 81(1):297-305. < PMID: 17056762 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com