

TECHNICAL DATA SHEET

CyFlow™ CD18 APC **Anti-Hu; Clone MEM-148**

REF CQ738281

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD18
Alternative Names	b2 integrin
Clone	MEM-148
Clonality	monoclonal
Format	APC
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	Pig
Quantity	100 tests
Immunogen	Peripheral blood mononuclear cells

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Specificity

The mouse monoclonal antibody MEM-148 recognizes an epitope on CD18 which is essentially inaccessible in intact integrin molecules on resting leukocytes, but is exposed on high-affinity state of LFA-1 or on unassociated CD18. CD18 is a 90-95 kDa type I transmembrane protein expressed on all leukocytes.

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD18 (integrin β2 subunit) forms heterodimers with four types of CD11 molecule to constitute leukocyte (β2) integrins: αLβ2 (CD11a/CD18, LFA-1), αMβ2 (CD11b/CD18, Mac-1, CR3), αXβ2 (CD11c/CD18) and αDβ2 (CD11d/CD18). In most cases, the response mediated by the integrin is a composite of the functions of its individual subunits. These integrins are essential for proper leukocyte migration, mediating intercellular contacts. Absence of CD18 leads to leukocyte adhesion deficiency-1; severe reduction of CD18 expression leads to the development of a psoriasiform skin disease. CD18 is also a target of Mannheimia (Pasteurella) haemolytica leukotoxin and is sufficient to mediate leukotoxin-mediated cytolysis.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

References

- Gao JX, Issekutz AC: Mac-1 (CD11b/CD108) is the predominant beta 2 (CD108) integrin mediating human neutrophil migration through synovial and dermal fibroblast barriers. *Immunology*. 1996 Jul; 88(3):463-70. < PMID: 8774366 >
- Kishimoto T, Goyert S, Kikutani H, Mason D, Miyasaka M, Moretta L, Ohno T, Okumura K, Shaw S, Springer TA, Sugamura K, Sugawara H, von dem Borne AEGK, Zola H (Eds): *Leucocyte Typing VI*. Garland Publishing Inc, New York. 1997; 1-1342. < NLM ID: 9712219 >
- Shang XZ, Issekutz AC: Contribution of CD11a/CD18, CD11b/CD18, ICAM-1 (CD54 and -2 (CD102 to human monocyte migration through endothelium and connective tissue fibroblast barriers. *Eur J Immunol*. 1998 Jun; 28(6):1970-9. < PMID: 9645379 >
- Drbal K, Angelisova P, Cerny J, Pavlistova D, Cebecauer M, Novak P, Horejsi V: Human leukocytes contain a large pool of free forms of CD18. *Biochem Biophys Res Commun*. 2000 Aug 28; 275(2):295-9. < PMID: 10964661 >
- Drbal K, Angelisova P, Cerny J, Hilgert I, Horejsi V: A novel anti-CD18 mAb recognizes an activation-related epitope and induces a high-affinity conformation in leukocyte integrins. *Immunobiology*. 2001 May; 203(4):687-98. < PMID: 11402502 >
- Drbal K, Angelisova P, Hilgert I, Cerny J, Novak P, Horejsi V: A proteolytically truncated form of free CD18, the common chain of leukocyte integrins, as a novel marker of activated myeloid cells. *Blood*. 2001 Sep 1; 98(5):1561-6. < PMID: 11520808 >
- Kess D, Peters T, Zamek J, Wickenhauser C, Tawadros S, Loser K, Varga G, Grabbe S, Nischt R, Sunderkötter C, Müller W, Krieg T, Scharffetter-Kochanek K: CD4+ T cell-associated pathophysiology critically depends on CD18 gene dose effects in a murine model of psoriasis. *J Immunol*. 2003 Dec 1; 171(11):5697-706. < PMID: 14634077 >
- Solovjov DA, Pluskota E, Plow EF: Distinct roles for the alpha and beta subunits in the functions of integrin alphaMbeta2. *J Biol Chem*. 2005 Jan 14; 280(2):1336-45. < PMID: 15485828 >
- Peters T, Sindrilaru A, Wang H, Oreshkova T, Renkl AC, Kess D, Scharffetter-Kochanek K: CD18 in monogenic and polygenic inflammatory processes of the skin. *J Invest Dermatol Symp Proc*. 2006 Sep; 11(1):42186. < PMID: 17069006 >
- Dassanayake RP, Maheswaran SK, Srikumaran S: Monomeric expression of bovine beta2-integrin subunits reveals their role in *Mannheimia haemolytica* leukotoxin-induced biological effects. *Infect Immun*. 2007 Oct; 75(10):5004-10. < PMID: 17698568 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Aubert M, Yoon M, Sloan DD, Spear PG, Jerome KR: The virological synapse facilitates herpes simplex virus entry into T cells. J Virol. 2009 Jun; 83(12):6171-83. < PMID: 19339346 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com