

TECHNICAL DATA SHEET

CyFlow™ CD18 Biotin **Anti-Hu; Clone MEM-48**

REF AT212653

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD18
Alternative Names	b2 integrin
Clone	MEM-48
Clonality	monoclonal
Format	Biotin

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	Dog
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Leukocytes of a patient suffering from LGL-type Leukemia

Specificity

The mouse monoclonal antibody MEM-48 recognizes an epitope involving residues 534-546 in cysteine-rich repeat 3 of the CD18 antigen. CD18 is a 90-95 kDa type I transmembrane protein expressed on all leukocytes.

Application

The reagent is designed for indirect immunofluorescence analysis by Flow Cytometry. Suggested working usage is 10 µg/ml. Indicated dilution is recommended starting point for use of this product, but working concentrations should be validated by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in Tris buffered saline (TBS) solution, pH ≈8.0, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD18 (integrin β2 subunit) forms heterodimers with four types of CD11 molecule to constitute leukocyte (β2) integrins: αLβ2 (CD11a/CD18, LFA-1), αMβ2 (CD11b/CD18, Mac-1, CR3), αXβ2 (CD11c/CD18) and αDβ2 (CD11d/CD18). In most cases, the response mediated by the integrin is a composite of the functions of its individual subunits. These integrins are essential for proper leukocyte migration, mediating intercellular contacts. Absence of CD18 leads to leukocyte adhesion deficiency-1; severe reduction of CD18 expression leads to the development of a psoriasiform skin disease. CD18 is also a target of

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Mannheimia (Pasteurella) haemolytica leukotoxin and is sufficient to mediate leukotoxin-mediated cytolysis.

References

- Persson H, Hennighausen L, Taub R, DeGrado W, Leder P: Antibodies to human c-myc oncogene product: evidence of an evolutionarily conserved protein induced during cell proliferation. Science. 1984 Aug 17; 225(4663):687-93. < PMID: 6431612 >
- Bazil V, Stefanova I, Hilgert I, Kristofova H, Vanek S, Horejsi V: Monoclonal antibodies against human leucocyte antigens (IV): Antibodies against subunits of the LFA-1 (CD11a/CD18) leucocyte-adhesion glycoprotein. Folia Biol (Praha). 1990; 36(1):41-50. < PMID: 1971601 >
- Larson RS, Springer TA: Structure and function of leukocyte integrins. Immunol Rev. 1990 Apr; 114:181-217. < PMID: 2196220 >
- Stefanova I, Horejsi V: Association of the CD59 and CD55 cell surface glycoproteins with other membrane molecules. J Immunol. 1991 Sep 1; 147(5):1587-92. < PMID: 1715364 >
- Schiff DE, Rae J, Martin TR, Davis BH, Curnutte JT: Increased phagocyte Fc gammaRI expression and improved Fc gamma-receptor-mediated phagocytosis after in vivo recombinant human interferon-gamma treatment of normal human subjects. Blood. 1997 Oct 15; 90(8):3187-94. < PMID: 9376602 >
- Gannon GA, Rhind SG, Suzui M, Zamecnik J, Sabiston BH, Shek PN, Shephard RJ: beta-Endorphin and natural killer cell cytolytic activity during prolonged exercise: Is there a connection?. Am J Physiol. 1998 Dec; 275(6-2):R1725-34. < PMID: 9843861 >
- Ottonello L, Epstein AL, Dapino P, Barbera P, Morone P, Dallegri F: Monoclonal Lym-1 antibody-dependent cytolysis by neutrophils exposed to granulocyte-macrophage colony-stimulating factor: intervention of Fc gammaRII (CD32), CD11b-CD18 integrins, and CD66b glycoproteins. Blood. 1999 May 15; 93(10):3505-11. < PMID: 10233903 >
- Garnotel R, Rittié L, Poitevin S, Monboisse JC, Nguyen P, Potron G, Maquart FX, Randoux A, Gillery P: Human blood monocytes interact with type I collagen through alpha x beta 2 integrin (CD11c-CD18, gp150-95). J Immunol. 2000 Jun 1; 164(11):5928-34. < PMID: 10820275 >
- Solovey AN, Gui L, Chang L, Enenstein J, Browne PV, Hebbel RP: Identification and functional assessment of endothelial P1H12. J Lab Clin Med. 2001 Nov; 138(5):322-31. < PMID: 11709656 >
- Dao MA, Arevalo J, Nolte JA: Reversibility of CD34 expression on human hematopoietic stem cells that retain the capacity for secondary reconstitution. Blood. 2003 Jan 1; 101(1):112-8. < PMID: 12589631 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Kerstan A, Hünig T: Cutting edge: distinct TCR- and CD28-derived signals regulate CD95L, Bcl-xL, and the survival of primary T cells. J Immunol. 2004 Feb 1; 172(3):1341-5. < PMID: 14734708 >
- Shan M, Klasse PJ, Banerjee K, Dey AK, Iyer SP, Dionisio R, Charles D, Campbell-Gardener L, Olson WC, Sanders RW, Moore JP: HIV-1 gp120 mannoses induce immunosuppressive responses from dendritic cells. PLoS Pathog. 2007 Nov; 3(11):e169. < PMID: 17983270 >
- Kuttruff S, Koch S, Kelp A, Pawelec G, Rammensee HG, Steinle A: NKp80 defines and stimulates a reactive subset of CD8 T cells. Blood. 2009 Jan 8; 113(2):358-69. < PMID: 18922855 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com