

TECHNICAL DATA SHEET

CyFlow™ CD18 FITC Anti-Ms; Clone M18/2

REF AQ070678

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD18
Alternative Names	b2 integrin
Clone	M18/2
Clonality	monoclonal
Format	FITC
Host / Isotype	Rat / IgG2a
Species Reactivity	Mouse
Negative Species Reactivity	Dog
Quantity [Concentration]	0.1 mg [0.5 mg/ml]
Immunogen	Murine cytotoxic T cell glycoproteins

Specificity

The rat monoclonal antibody M18/2 recognizes mouse CD18 antigen, a 95 kDa type I transmembrane protein expressed on all leukocytes.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Suggested working usage is 3 µg/ml. Indicated dilution is recommended starting point for use of this product, but working concentrations should be validated by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD18 (integrin β2 subunit) forms heterodimers with four types of CD11 molecule to constitute leukocyte (β2) integrins: αLβ2 (CD11a/CD18, LFA-1), αMβ2 (CD11b/CD18, Mac-1, CR3), αXβ2 (CD11c/CD18) and αDβ2 (CD11d/CD18). In most cases, the response mediated by the integrin is a composite of the functions of its individual subunits. These integrins are essential for proper leukocyte migration, mediating intercellular contacts.

References

- Sanchez-Madrid F, Simon P, Thompson S, Springer TA: Mapping of antigenic and functional epitopes on the alpha- and beta-subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. J Exp Med. 1983 Aug 1; 158(2):586-602. < PMID: 6193226 >
- Zahalka MA, Okon E, Naor D: Blocking lymphoma invasiveness with a monoclonal antibody directed against the beta-chain of the leukocyte adhesion molecule (CD18). J Immunol. 1993 May 15; 150(10):4466-77. < PMID: 8097756 >
- Zahalka MA, Naor D: Beta 2-integrin dependent aggregate formation between LB T cell lymphoma and spleen cells: assessment of correlation with spleen invasiveness. Int Immunol. 1994 Jun; 6(6):917-24. < PMID: 7522049 >
- Avni O, Pur Z, Yefenof E, Baniyash M: Complement receptor 3 of macrophages is associated with galectin-1-like protein. J Immunol. 1998 Jun 15; 160(12):6151-8. < PMID: 9637533 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Abraham C, Miller J: Molecular mechanisms of IL-2 gene regulation following costimulation through LFA-1. J Immunol. 2001 Nov 1; 167(9):5193-201. < PMID: 11673532 >
- Sakurai E, Taguchi H, Anand A, Ambati BK, Gragoudas ES, Miller JW, Adamis AP, Ambati J: Targeted disruption of the CD18 or ICAM-1 gene inhibits choroidal neovascularization. Invest Ophthalmol Vis Sci. 2003 Jun; 44(6):2743-9. < PMID: 12766082 >
- Barlow SC, Langston W, Matthews KM, Chidlow JH Jr, Kevil CG: CD18 deficiency protects against multiple low-dose streptozotocin-induced diabetes. Am J Pathol. 2004 Dec; 165(6):1849-52. < PMID: 15579429 >
- Watts GM, Beurskens FJ, Martin-Padura I, Ballantyne CM, Klickstein LB, Brenner MB, Lee DM: Manifestations of inflammatory arthritis are critically dependent on LFA-1. J Immunol. 2005 Mar 15; 174(6):3668-75. < PMID: 15749905 >
- Varga G, Balkow S, Wild MK, Stadtbauer A, Krummen M, Rothoeft T, Higuchi T, Beissert S, Wethmar K, Scharffetter-Kochanek K, Vestweber D, Grabbe S: Active MAC-1 (CD11b/CD18) on DCs inhibits full T-cell activation. Blood. 2007 Jan 15; 109(2):661-9. < PMID: 17003381 >
- Cullere X, Lauterbach M, Tsuboi N, Mayadas TN: Neutrophil-selective CD18 silencing using RNA interference in vivo. Blood. 2008 Apr 1; 111(7):3591-8. < PMID: 18216298 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com