

TECHNICAL DATA SHEET

CyFlow™ CD18 Purified Anti-Ms; Clone M18/2

REF CK708893

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD18
Alternative Names	b2 integrin
Clone	M18/2
Clonality	monoclonal
Format	Purified
Host / Isotype	Rat / IgG2a
Species Reactivity	Mouse
Negative Species Reactivity	Dog
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Murine cytotoxic T cell glycoproteins

Specificity

The rat monoclonal antibody M18/2 recognizes mouse CD18 antigen, a 95 kDa type I transmembrane protein expressed on all leukocytes.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Western blot
- Immunohistochemistry
- Functional assays

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD18 (integrin β 2 subunit) forms heterodimers with four types of CD11 molecule to constitute leukocyte (β 2) integrins: α L β 2 (CD11a/CD18, LFA-1), α M β 2 (CD11b/CD18, Mac-1, CR3), α X β 2 (CD11c/CD18) and α D β 2 (CD11d/CD18). In most cases, the response mediated by the integrin is a composite of the functions of its individual subunits. These integrins are essential for proper leukocyte migration, mediating intercellular contacts.

References

- Sanchez-Madrid F, Simon P, Thompson S, Springer TA: Mapping of antigenic and functional epitopes on the alpha- and beta-subunits of two related mouse glycoproteins involved in cell interactions, LFA-1 and Mac-1. J Exp Med. 1983 Aug 1; 158(2):586-602. < PMID: 6193226 >
- Zahalka MA, Okon E, Naor D: Blocking lymphoma invasiveness with a monoclonal antibody directed against the beta-chain of the leukocyte adhesion molecule (CD18). J Immunol. 1993 May 15; 150(10):4466-77. < PMID: 8097756 >
- Zahalka MA, Naor D: Beta 2-integrin dependent aggregate formation between LB T cell lymphoma and spleen cells: assessment of correlation with spleen invasiveness. Int Immunol. 1994 Jun; 6(6):917-24. < PMID: 7522049 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Avni O, Pur Z, Yefenof E, Baniyash M: Complement receptor 3 of macrophages is associated with galectin-1-like protein. J Immunol. 1998 Jun 15; 160(12):6151-8. < PMID: 9637533 >
- Abraham C, Miller J: Molecular mechanisms of IL-2 gene regulation following costimulation through LFA-1. J Immunol. 2001 Nov 1; 167(9):5193-201. < PMID: 11673532 >
- Sakurai E, Taguchi H, Anand A, Ambati BK, Gragoudas ES, Miller JW, Adamis AP, Ambati J: Targeted disruption of the CD18 or ICAM-1 gene inhibits choroidal neovascularization. Invest Ophthalmol Vis Sci. 2003 Jun; 44(6):2743-9. < PMID: 12766082 >
- Barlow SC, Langston W, Matthews KM, Chidlow JH Jr, Kevil CG: CD18 deficiency protects against multiple low-dose streptozotocin-induced diabetes. Am J Pathol. 2004 Dec; 165(6):1849-52. < PMID: 15579429 >
- Watts GM, Beurskens FJ, Martin-Padura I, Ballantyne CM, Klickstein LB, Brenner MB, Lee DM: Manifestations of inflammatory arthritis are critically dependent on LFA-1. J Immunol. 2005 Mar 15; 174(6):3668-75. < PMID: 15749905 >
- Varga G, Balkow S, Wild MK, Stadtbauer A, Krummen M, Rothoeft T, Higuchi T, Beissert S, Wethmar K, Scharffetter-Kochanek K, Vestweber D, Grabbe S: Active MAC-1 (CD11b/CD18) on DCs inhibits full T-cell activation. Blood. 2007 Jan 15; 109(2):661-9. < PMID: 17003381 >
- Cullere X, Lauterbach M, Tsuboi N, Mayadas TN: Neutrophil-selective CD18 silencing using RNA interference in vivo. Blood. 2008 Apr 1; 111(7):3591-8. < PMID: 18216298 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com