

TECHNICAL DATA SHEET

CyFlow™ CD1a APC Anti-Hu; Clone SK9

REF CM362473

**For Research Use Only.
Not for use in diagnostic or therapeutic procedures.**

Specifications

Antigen	CD1a
Alternative Names	R4, HTA1, CD1
Clone	SK9
Clonality	monoclonal
Format	APC
Host / Isotype	Mouse / IgG2b
Species Reactivity	Human
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Human thymocytes

Specificity

The mouse monoclonal antibody SK9 recognizes CD1a antigen, a 49 kDa polypeptide associated with β 2-microglobulin expressed on cortical thymocytes (strongly), Langerhans cells, dendritic cells and some T cell leukemias and lymphomas.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD1a, together with CD1b and CD1c, belongs to group 1 of CD1 glycoproteins. These proteins serve as antigen-presenting molecules for a subset of T cells that responds to specific lipids and glycolipids found in the cell walls of bacterial pathogens or self-glycolipid antigens such as gangliosides, and they have also roles in antiviral immunity. Unlike CD1b, CD1a is excluded from late endosomal compartments and instead traffics independently in the recycling pathway of the early endocytic system, and CD1a antigen presentation is independent upon vesicular acidification.

References

- Wood GS, Burns BF, Dorfman RF, Warnke RA: The immunohistology of non-T cells in the acquired immunodeficiency syndrome. *Am J Pathol.* 1985 Sep; 120(3):371-9. < PMID: 3898858 >
- Zimmer MI, Larregina AT, Castillo CM, Capuano S 3rd, Falo LD Jr, Murphey-Corb M, Reinhart TA, Barratt-Boyes SM: Disrupted homeostasis of Langerhans cells and interdigitating dendritic cells in monkeys with AIDS. *Blood.* 2002 Apr 15; 99(8):2859-68. < PMID: 11929776 >
- Tenca C, Merlo A, Zarccone D, Saverino D, Bruno S, De Santanna A, Ramarli D, Fabbi M, Pesce C, Deaglio S, Ciccone E, Malavasi F, Grossi CE: Death of T cell precursors in the human thymus: a role for CD38. *Int Immunol.* 2003 Sep; 15(9):1105-16. < PMID: 12917263 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Brown KN, Trichel A, Barratt-Boyes SM: Parallel loss of myeloid and plasmacytoid dendritic cells from blood and lymphoid tissue in simian AIDS. J Immunol. 2007 Jun 1; 178(11):6958-67. < PMID: 17513745 >
- Awasthi S, Wolf R, White G: Ontogeny and phagocytic function of baboon lung dendritic cells. Immunol Cell Biol. 2009 Jul; 87(5):419-27. < PMID: 19223853 >
- Kader M, Smith AP, Guiducci C, Wonderlich ER, Normolle D, Watkins SC, Barrat FJ, Barratt-Boyes SM: Blocking TLR7- and TLR9-mediated IFN- α : production by plasmacytoid dendritic cells does not diminish immune activation in early SIV infection. PLoS Pathog. 2013; 9(7):e1003530. < PMID: 23935491 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com