

TECHNICAL DATA SHEET

CyFlow™ CD264 PE Anti-Hu; Clone TRAIL-R4-01

REF CY292936

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD264
Alternative Names	TRAILR4, DCR2, TRUNDD, TNFRSF10d
Clone	TRAIL-R4-01
Clonality	monoclonal
Format	PE
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [0.1 mg/ml]
Immunogen	TRAIL-R4 (aa 1-210) - hlgGhc fusion protein

Specificity

The mouse monoclonal antibody TRAIL-R4-01 recognizes CD264 (TRAIL-R4) antigen, a 42 kDa transmembrane protein expressed on various blood cells.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Suggested working usage is 3 µg/ml. Indicated dilution is recommended starting point for use of this product, but working concentrations should be validated by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD264 (TRAIL-R4, TR4, DcR2, TRUNDD), expressed mainly on CD8+ and NK cells, belongs to receptors of TRAIL, a TNF-like membrane toxic protein that induces apoptosis in many tumor cells, but not in normal cells. TRAIL-R4, however, contains partially truncated death domain, thus it is unable to induce apoptosis and serves as a negative regulator of apoptotic signaling by impairment death-inducing signaling complex (DISC) processing. TRAIL-R4 interacts with death receptor 5 (DR5) in the native DISC in a TRAIL-dependent manner and prevents its corecruitment with death receptor 4 (DR4).

References

- Clancy L, Mruk K, Archer K, Woelfel M, Mongkolsapaya J, Screaton G, Lenardo MJ, Chan FK: Preligand assembly domain-mediated ligand-independent association between TRAIL receptor 4 (TR4) and TR2 regulates TRAIL-induced apoptosis. Proc Natl Acad Sci USA. 2005 Dec 13; 102(50):18099-104. < PMID: 16319225 >
- Sanlioglu AD, Dirice E, Aydin C, Erin N, Koksoy S, Sanlioglu S: Surface TRAIL decoy receptor-4 expression is correlated with TRAIL resistance in MCF7 breast cancer cells. BMC Cancer. 2005 May 25; 5(1):54. < PMID: 18518976 >
- Mérino D, Lalaoui N, Morizot A, Schneider P, Solary E, Micheau O: Differential inhibition of TRAIL-mediated DR5-DISC formation by decoy receptors 1 and 2. Mol Cell Biol. 2006 Oct; 26(19):7046-55. < PMID: 16980609 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Deligezer U, Dalay N: Expression of the TRAIL Receptors in Blood Mononuclear Cells in Leukemia. Pathol Oncol Res. 2007; 13(4):290-4. < PMID: 18158563 >
- Falschlehner C, Emmerich CH, Gerlach B, Walczak H: TRAIL signalling. Int J Biochem Cell Biol. 2007; 39(7-8):1462-75. < PMID: 17403612 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com