

TECHNICAL DATA SHEET

CyFlow™ CD29 FITC Anti-Hu; Clone MEM-101A

REF BZ181097

**For Research Use Only.
Not for use in diagnostic or therapeutic procedures.**

Specifications

Antigen	CD29
Alternative Names	Integrin β 1
Clone	MEM-101A
Clonality	monoclonal
Format	FITC
Host / Isotype	Mouse / IgG1
Species Reactivity	Human, Pig Dog
Negative Species Reactivity	Mouse
Quantity	100 tests
Immunogen	Raji cells: human Burkitt's lymphoma cell line

Specificity

The mouse monoclonal antibody MEM-101A recognizes CD29 antigen, a 130 kDa single chain type I glycoprotein expressed as a heterodimer (non-covalently associated with the integrin α subunits 1-6).

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

CD29 is broadly expressed on majority of hematopoietic and non-hematopoietic cells (leukocytes, platelets, fibroblasts, endothelial cells, epithelial cells and mast cells).

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD29 (β1 integrin subunit, GPIIa) forms non-covalently linked heterodimers with at least 6 different α chains (α1-α6, CD49a-f) determining the binding properties of β1 (VLA) integrins. These integrins mediate cell adhesion to collagen, fibronectin, laminin and other extracellular matrix (ECM) components. This interaction hinders cell death, whereas disruption of anchorage to ECM leads to apoptosis. Decreased expression of most β1 integrins correlates with acquiring multidrug resistance of tumor cells during selection in presence of antitumor drug. In platelets, translocation of intracellular pool of β1 integrins to the plasma membrane following thrombin stimulation. These integrins are also up-regulated in leukocytes during emigration and extravascular migration and appear to be critically involved in regulating the immune cell trafficking from blood to tissue, as well as in regulating tissue damage and disease symptoms related to inflammatory bowel disease. Through a β1 integrin-dependent mechanism, fibronectin and type I collagen enhance cytokine secretion of human airway smooth muscle in response to IL-1β.

References

- Wencel-Drake JD, Dieter MG, Lam SC: Immunolocalization of beta 1 integrins in platelets and platelet-derived microvesicles. Blood. 1993 Aug 15; 82(4):1197-203. < PMID: 7688992 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Kishimoto T, Goyert S, Kikutani H, Mason D, Miyasaka M, Moretta L, Ohno T, Okumura K, Shaw S, Springer TA, Sugamura K, Sugawara H, von dem Borne AEGK, Zola H (Eds): Leucocyte Typing VI. Garland Publishing Inc, New York. 1997; 1-1342. < NLM ID: 9712219 >
- Werr J, Johansson J, Eriksson EE, Hedqvist P, Ruoslahti E, Lindbom L: Integrin alpha(2)beta(1) (VLA-2) is a principal receptor used by neutrophils for locomotion in extravascular tissue. Blood. 2000 Mar 1; 95(5):1804-9. < PMID: 10688841 >
- Peng Q, Lai D, Nguyen TT, Chan V, Matsuda T, Hirst SJ: Multiple beta 1 integrins mediate enhancement of human airway smooth muscle cytokine secretion by fibronectin and type I collagen. J Immunol. 2005 Feb 15; 174(4):2258-64. < PMID: 15699160 >
- Lundberg S, Lindholm J, Lindbom L, Hellström PM, Werr J: Integrin alpha2beta1 regulates neutrophil recruitment and inflammatory activity in experimental colitis in mice. Inflamm Bowel Dis. 2006 Mar; 12(3):172-7. < PMID: 16534417 >
- Morozevich GE, Kozlova NI, Preobrazhenskaya ME, Ushakova NA, Eltsov IA, Shtil AA, Berman AE: The role of beta1 integrin subfamily in anchorage-dependent apoptosis of breast carcinoma cells differing in multidrug resistance. Biochemistry (Mosc). 2006 May; 71(5):489-95. < PMID: 16732726 >
- Simova S, Klima M, Cermak L, Sourkova V, Andera L: Arf and Rho GAP adapter protein ARAP1 participates in the mobilization of TRAIL-R1/DR4 to the plasma membrane. Apoptosis. 2008 Mar; 13(3):423-36. < PMID: 18165900 >
- Plánka L, Necas A, Srnec R, Rauser P, Starý D, Jancár J, Amler E, Filová E, Hlucilová J, Kren L, Gál P: Use of allogenic stem cells for the prevention of bone bridge formation in miniature pigs. Physiol Res. 2009; 58(6):885-93. < PMID: 19093735 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com