

TECHNICAL DATA SHEET

CyFlow™ CD3 FITC **Anti-Ms; Clone 145-2C11**

REF AJ219456

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD3
Alternative Names	T3, Leu4
Clone	145-2C11
Clonality	monoclonal
Format	FITC
Host / Isotype	Hamster / IgG
Species Reactivity	Mouse
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [0.5 mg/ml]
Immunogen	Mouse BM10-37 cytotoxic T lymphocytes

Specificity

The Armenian hamster monoclonal antibody 145-2C11 recognizes mouse CD3 (ε subunit). This antibody is commonly used as a phenotypic marker for mouse T cells.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Suggested working usage is 2 µg/ml. Indicated dilution is recommended starting point for use of this product, but working concentrations should be validated by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3 subunits CD3γ, CD3 δ, CD3ε and CD3ζ. These CD3 subunits are structurally related members of the immunoglobulins superfamily encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3 may play a role in TCR-induced growth arrest, cell survival and proliferation.

References

- Leo O, Foo M, Sachs DH, Samelson LE, Bluestone JA: Identification of a monoclonal antibody specific for a murine T3 polypeptide. Proc Natl Acad Sci USA. 1987 Mar; 84(5):1374-8. < PMID: 2950524 >
- Samelson LE, O'Shea JJ, Luong H, Ross P, Urdahl KB, Klausner RD, Bluestone J: T cell antigen receptor phosphorylation induced by an anti-receptor antibody. J Immunol. 1987 Oct 15; 139(8):2708-14. < PMID: 2443570 >
- Payer E, Elbe A, Stingl G: Circulating CD3+/T cell receptor V gamma 3+ fetal murine thymocytes home to the skin and give rise to proliferating dendritic epidermal T cells. J Immunol. 1991 Apr 15; 146(8):2536-43. < PMID: 1826696 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Salvadori S, Gansbacher B, Pizzimenti AM, Zier KS: Abnormal signal transduction by T cells of mice with parental tumors is not seen in mice bearing IL-2-secreting tumors. J Immunol. 1994 Dec 1; 153(11):5176-82. < PMID: 7963575 >
- Jacobs H, Vandeputte D, Tolkamp L, de Vries E, Borst J, Berns A: CD3 components at the surface of pro-T cells can mediate pre-T cell development in vivo. Eur J Immunol. 1994 Apr; 24(4):934-9. < PMID: 8149963 >
- Vossen AC, Tibbe GJ, Kroos MJ, van de Winkel JG, Benner R, Savelkoul HF: Fc receptor binding of anti-CD3 monoclonal antibodies is not essential for immunosuppression, but triggers cytokine-related side effects. Eur J Immunol. 1995 Jun; 25(6):1492-6. < PMID: 7614975 >
- Henrickson M, Reid J, Bellet JS, Sawchuk SS, Hirsch R: Comparison of in vivo efficacy and mechanism of action of antimurine monoclonal antibodies directed against TCR alpha beta (H57-597) and CD3 (145-2C11). Transplantation. 1995 Oct 27; 60(8):828-35. < PMID: 7482743 >
- Kinnaert P, Pradier O, Bournonville B, Habrant C, Goldman M, Van Geertruyden N: Role of CD18-dependent and CD18-independent mechanisms in the increased leukocyte adhesiveness and in the variations of circulating white blood cell populations induced by anti-CD3 monoclonal antibodies. Transpl Int. 1996; 9(4):386-91. < PMID: 8819275 >
- Kearse KP: Calnexin associates with monomeric and oligomeric (disulfide-linked) CD3delta proteins in murine T lymphocytes. J Biol Chem. 1998 Jun 5; 273(23):14152-7. < PMID: 9603915 >
- Han WR, Murray-Segal LJ, Gershenson A, Zhang JG, Hodder AN, Pietersz GA, Mottram PL: Idarubicin-145-2C11-F(ab')₂ promotes peripheral tolerance and reduces chronic vascular disease in mouse cardiac allografts. Transpl Immunol. 1999 Dec; 7(4):207-13. < PMID: 10638833 >
- Tilley SL, Jaradat M, Stapleton C, Dixon D, Hua X, Erikson CJ, McCaskill JG, Chason KD, Liao G, Jania L, Koller BH, Jetten AM: Retinoid-related orphan receptor gamma controls immunoglobulin production and Th1/Th2 cytokine balance in the adaptive immune response to allergen. J Immunol. 2007 Mar 1; 178(5):3208-18. < PMID: 17312169 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com