

TECHNICAL DATA SHEET

CyFlow™ CD3 Purified Anti-Hu; Clone MEM-57

REF AL894369

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD3
Alternative Names	T3, Leu4
Clone	MEM-57
Clonality	monoclonal
Format	Purified
Host / Isotype	Mouse / IgG2a
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Human thymocytes and T lymphocytes

Specificity

The mouse monoclonal antibody MEM-57 recognizes γ - ϵ and δ - ϵ dimers of human CD3 complex, a part of a bigger multisubunit T cell receptor complex (CD3/TCR) expressed on peripheral blood T lymphocytes and mature thymocytes.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Functional assays

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD3 complex is crucial in transducing antigen-recognition signals into the cytoplasm of T cells and in regulating the cell surface expression of the TCR complex. T cell activation through the antigen receptor (TCR) involves the cytoplasmic tails of the CD3 subunits CD3 γ , CD3 δ , CD3 ϵ and CD3 ζ . These CD3 subunits are structurally related members of the immunoglobulins super family encoded by closely linked genes on human chromosome 11. The CD3 components have long cytoplasmic tails that associate with cytoplasmic signal transduction molecules. This association is mediated at least in part by a double tyrosine-based motif present in a single copy in the CD3 subunits. CD3 may play a role in TCR-induced growth arrest, cell survival and proliferation. The CD3 antigen is present on 68-82% of normal peripheral blood lymphocytes, 65-85% of thymocytes and Purkinje cells in the cerebellum. It is never expressed on B or NK cells. Decreased percentages of T lymphocytes may be observed in some autoimmune diseases.

References

- McMichael AJ, Beverley PCL, Cobbold S, et al. (Eds): Leucocyte Typing III, White Cell Differentiation Antigens. Oxford University Press, Oxford. 1987; 1-1050. < NLM ID: 8913266 >
- Horejsí V, Angelisová P, Bazil V, Kristofová H, Stoyanov S, Stefanová I, Hausner P, Vosecký M, Hilgert I: Monoclonal antibodies against human leucocyte antigens (II); Antibodies against CD45 (T200), CD3 (T3), CD43, CD10 (CALLA), transferrin receptor (T9), a novel broadly expressed 18-kDa antigen

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- (MEM-43) and a novel antigen of restricted expression (MEM-74). *Folia Biol (Praha)*. 1988; 34(1):23-34. < PMID: 2968928 >
- Knapp W, Dorken B, Gilks W, Rieber EP, Schmidt RE, Stein H, von dem Borne AEGK (Eds): *Leucocyte Typing IV*. Oxford University Press, Oxford. 1989; 1-1820. < NLM ID: 8914679 >
 - Soucek J, Chudomel V, Hrubá A, Lindnerová G: Induction of NK and LAK activities in human lymphocyte culture by a cytosol fraction from leukemic myeloblasts and by monoclonal antibody CD 3. *Neoplasma*. 1991; 38(1):33-41. < PMID: 2011208 >
 - Hilgert I, Franěk F, Stefanová I, Kaslík J, Jirka J, Kristofová H, Rossmann P, Soucek J, Horejsi V: Therapeutic in vivo use of the A1-CD3 monoclonal antibody. *Transplantation*. 1993; 55:435. < PMID: 8434399 >
 - Soucek J, Hilgert I, Budová I, Lindnerová G: Augmentation of NK cell activity and proliferation in cultured lymphocytes of leukemic patients by monoclonal antibodies CD3 and interleukin-2. *Neoplasma*. 1994; 41(2):75-81. < PMID: 8208318 >
 - Dave VP, Cao Z, Browne C, Alarcon B, Fernandez-Miguel G, Lafaille J, de la Hera A, Tonegawa S, Kappes DJ: CD3 delta deficiency arrests development of the alpha beta but not the gamma delta T cell lineage. *EMBO J*. 1997 Mar 17; 16(6):1360-70. < PMID: 9135151 >
 - Panyi G, Bagdány M, Bodnár A, Vámosi G, Szentesi G, Jenei A, Mátyus L, Varga S, Waldmann TA, Gáspár R, Damjanovich S: Colocalization and nonrandom distribution of Kv1.3 potassium channels and CD3 molecules in the plasma membrane of human T lymphocytes. *Proc Natl Acad Sci USA*. 2003 Mar 4; 100(5):2592-7. < PMID: 12604782 >
 - Brdicková N, Brdicka T, Angelisová P, Horváth O, Spicka J, Hilgert I, Paces J, Simeoni L, Kliche S, Merten C, Schraven B, Horejsi V: LIME: a new membrane Raft-associated adaptor protein involved in CD4 and CD8 coreceptor signaling. *J Exp Med*. 2003 Nov 17; 198(10):1453-62. < PMID: 14610046 >
 - Huang Y, Wange RL: T cell receptor signaling: beyond complex complexes. *J Biol Chem*. 2004 Jul 9; 279(28):28827-3. < PMID: 15084594 >
 - Kuhns MS, Davis MM, Garcia KC: Deconstructing the form and function of the TCR/CD3 complex. *Immunity*. 2006 Feb; 24(2):133-9. < PMID: 16473826 >
 - Alarcón B, Swamy M, van Santen HM, Schamel WW: T-cell antigen-receptor stoichiometry: pre-clustering for sensitivity. *EMBO Rep*. 2006 May; 7(5):490-5. < PMID: 16670682 >
 - Drbal K, Moertelmaier M, Holzhauser C, Muhammad A, Fuerthbauer E, Howorka S, Hinterberger M, Stockinger H, Schütz GJ: Single-molecule microscopy reveals heterogeneous dynamics of lipid raft components upon TCR engagement. *Int Immunol*. 2007 May; 19(5):675-84. < PMID: 17446208 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
 Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com