

TECHNICAL DATA SHEET

CyFlow™ CD305 PerCP-Cy5.5 Anti-Hu; Clone NKTA255

REF CT614380

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD305
Alternative Names	LAIR1, LAIR-1
Clone	NKTA255
Clonality	monoclonal
Format	PerCP-Cy5.5
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Activated NK cells and CD3- thymocytes

Specificity

The mouse monoclonal antibody NKTA255 recognizes CD305 (LAIR1) antigen, a 40 kDa type I transmembrane glycoprotein expressed on NK, T, and B cells, monocytes, dendritic cells, eosinophils, basophils, mast cells, CD34+ hematopoietic progenitor cells and thymocytes.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD305 (LAIR1; leukocyte-associated Ig-like receptor 1) is an inhibitory receptor found on many types of peripheral blood cells. It serves to suppress cell cytotoxicity, activation, proliferation, and differentiation regarding autoantigens via its two intracellular ITIM sites. CD305 belongs to the immunoglobulin superfamily and the leukocyte-associated inhibitory receptor family of proteins. It reacts with collagen ligands.

References

- Poggi A, Pella N, Morelli L, Spada F, Revello V, Sivori S, Augugliaro R, Moretta L, Moretta A: p40, a novel surface molecule involved in the regulation of the non-major histocompatibility complex-restricted cytolytic activity in humans. Eur J Immunol. 1995 Feb; 25(2):369-76. < PMID: 7875198 >
- Poggi A, Tomasello E, Revello V, Nanni L, Costa P, Moretta L: p40 molecule regulates NK cell activation mediated by NK receptors for HLA class I antigens and TCR-mediated triggering of T lymphocytes. Int Immunol. 1997 Sep; 9(9):1271-9. < PMID: 9310830 >
- Poggi A, Tomasello E, Ferrero E, Zocchi MR, Moretta L: p40/LAIR-1 regulates the differentiation of peripheral blood precursors to dendritic cells induced by granulocyte-monocyte colony-stimulating factor. Eur J Immunol. 1998 Jul; 28(7):2086-91. < PMID: 9692876 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Warren HS: The Eighth Human Leucocyte Differentiation Antigen (HLDA8) Workshop: natural killer cell section report. Cell Immunol. 2005 Jul-Aug; 236(1-2):17-20. < PMID: 16157321 >
- Meyaard L: The inhibitory collagen receptor LAIR-1 (CD305). J Leukoc Biol. 2008 Apr; 83(4):799-803. < PMID: 18063695 >
- Colombo BM, Canevali P, Magnani O, Rossi E, Puppo F, Zocchi MR, Poggi A: Defective expression and function of the leukocyte associated Ig-like receptor 1 in B lymphocytes from systemic lupus erythematosus patients. PLoS One. 2012; 7(2):e31903. < PMID: 22355402 >
- Son M, Santiago-Schwarz F, Al-Abed Y, Diamond B: C1q limits dendritic cell differentiation and activation by engaging LAIR-1. Proc Natl Acad Sci USA. 2012 Nov 13; 109(46):E3160-7. < PMID: 23093673 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Cy and CyDye are trademarks of GE Healthcare.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com