

TECHNICAL DATA SHEET

CyFlow™ CD44 Biotin Anti-Hu; Clone MEM-85

REF CQ603573

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD44
Alternative Names	H-CAM, Pgp-1
Clone	MEM-85
Clonality	monoclonal
Format	Biotin
Host / Isotype	Mouse / IgG2b
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Leukocytes of a patient suffering from LGL-type Leukemia

Specificity

The mouse monoclonal antibody MEM-85 recognizes both cell surface-expressed and soluble form of CD44 antigen, a 80-95 kDa transmembrane glycoprotein (hyaladherin family) present on the most of cells

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

and tissues (leukocytes, endothelial cells, mesenchymal cells, etc.); it is negative on platelets and hepatocytes.

Application

The reagent is designed for indirect immunofluorescence analysis by Flow Cytometry. Suggested working usage is 1 µg/ml. Indicated dilution is recommended starting point for use of this product, but working concentrations should be validated by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD44 is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculation and homing, it can thus serve e.g. as a modulator of macrophage recruitment in response to pathogen. Although CD44 functions are essential for physiological activities of normal cells, elevated CD44 expression correlates with poor prognosis in many carcinomas, facilitating tumor growth and metastasis, antiapoptosis and directional motility of cancer cells.

References

- Stefanová I, Hilgert I, Bazil V, Kristofová H, Horejsí V: Human leucocyte surface glycoprotein CDw44 and lymphocyte homing receptor are identical molecules. Immunogenetics. 1989; 29(6):402-4. < PMID: 2659502 >
- Knapp W, Dorken B, Gilks W, Rieber EP, Schmidt RE, Stein H, von dem Borne AEGK (Eds): Leucocyte Typing IV. Oxford University Press, Oxford. 1989; 1-1820. < NLM ID: 8914679 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Bazil V, Strominger JL: Metalloprotease and serine protease are involved in cleavage of CD43, CD44, and CD16 from stimulated human granulocytes: Induction of cleavage of L-selectin via CD16. *J Immunol.* 1994 Feb 1; 152(3):1314-22. < PMID: 7507963 >
- Litvinov SV, Velders MP, Bakker HA, Fleuren GJ, Warnaar SO: Ep-CAM: a human epithelial antigen is a homophilic cell-cell adhesion molecule. *J Cell Biol.* 1994 Apr; 125(2):437-46. < PMID: 8163559 >
- Kishimoto T, Goyert S, Kikutani H, Mason D, Miyasaka M, Moretta L, Ohno T, Okumura K, Shaw S, Springer TA, Sugamura K, Sugawara H, von dem Borne AEGK, Zola H (Eds): *Leucocyte Typing VI*. Garland Publishing Inc, New York. 1997; 1-1342. < NLM ID: 9712219 >
- Schmidt CS, Liu J, Zhang T, Song HY, Sandusky G, Mintze K, Benschop RJ, Glasebrook A, Yang DD, Na S: Enhanced B cell expansion, survival, and humoral responses by targeting death receptor 6. *J Exp Med.* 2003 Jan 6; 197(1):51-62. < PMID: 12515813 >
- Suárez L, Vidriales MB, García-Laraña J, Sanz G, Moreno MJ, López A, Barrena S, Martínez R, Tormo M, Palomera L, Lavilla E, López-Berges MC, de Santiago M, de Equiza ME, Miguel JF, Orfao A: CD34+ cells from acute myeloid leukemia, myelodysplastic syndromes, and normal bone marrow display different apoptosis and drug resistance-associated phenotypes. *Clin Cancer Res.* 2004 Nov 15; 10(22):7599-606. < PMID: 15569991 >
- Holleman A, den Boer ML, Cheek MH, Kazemier KM, Pei D, Downing JR, Janka-Schaub GE, Göbel U, Graubner UB, Pui CH, Evans WE, Pieters R: Expression of the outcome predictor in acute leukemia 1 (OPAL1 gene is not an independent prognostic factor in patients treated according to COALL or St Jude protocols. *Blood.* 2006 Sep 15; 108(6):1984-90. < PMID: 16709928 >
- Subramaniam V, Gardner H, Jothy S: Soluble CD44 secretion contributes to the acquisition of aggressive tumor phenotype in human colon cancer cells. *Exp Mol Pathol.* 2007 Dec; 83(3):341-6. < PMID: 17945213 >
- Kolar GR, Mehta D, Pelayo R, Capra JD: A novel human B cell subpopulation representing the initial germinal center population to express AID. *Blood.* 2007 Mar 15; 109(6):2545-52. < PMID: 17132718 >
- Ide K, Wang H, Tahara H, Liu J, Wang X, Asahara T, Sykes M, Yang YG, Ohdan H: Role for CD47-SIRPalpha signaling in xenograft rejection by macrophages. *Proc Natl Acad Sci USA.* 2007 Mar 20; 104(12):5062-6. < PMID: 17360380 >
- Viel EC, Lemarié CA, Benkirane K, Paradis P, Schiffrin EL: Immune regulation and vascular inflammation in genetic hypertension. *Am J Physiol Heart Circ Physiol.* 2010 Mar; 298(3):H938-44. < PMID: 20044442 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
 Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com