

TECHNICAL DATA SHEET

CyFlow™ CD45R0 FITC Anti-Hu; Clone UCHL1

REF BA212505

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD45R0
Alternative Names	UCHL-1
Clone	UCHL1
Clonality	monoclonal
Format	FITC
Host / Isotype	Mouse / IgG2a
Species Reactivity	Human
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Human IL-2 dependent T cells

Specificity

The mouse monoclonal antibody UCHL1 recognizes CD45R0, a 180 kDa low molecular weight isoform of the leukocyte common antigen (LCA). CD45R0 is expressed on a subset of memory/activated T cells and on cortical thymocytes.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD45R0 is the shortest isoform of a receptor-type protein tyrosine phosphatase, CD45 glycoprotein. CD45 is crucial in lymphocyte development and antigen signaling, serving as an important regulator of Src-family kinases, promotes cell survival by modulating integrin-mediated signal transduction pathway and is also involved in DNA fragmentation during apoptosis. CD45 isoforms differ in their extracellular domains, whereas they share identical transmembrane and cytoplasmic domains. These isoforms differ in their ability to translocate into the glycosphingolipid-enriched membrane domains and their expression depends on cell type and physiological state of the cell. CD45R0 is expressed e.g. on macrophages, CD8+ T cells, activated T cells and myeloma cells.

References

- Norton AJ, Ramsay AD, Smith SH, Beverley PC, Isaacson PG: Monoclonal antibody (UCHL{01} that recognises normal and neoplastic T cells in routinely fixed tissues. J Clin Pathol. 1986 Apr; 39(4):399-405. < PMID: 2422219 >
- Smith SH, Brown MH, Rowe D, Callard RE, Beverley PC: Functional subsets of human helper-inducer cells defined by a new monoclonal antibody, UCHL1. Immunology. 1986 May; 58(1):63-70. < PMID: 2423439 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- McMichael AJ, Beverley PCL, Cobbold S, et al. (Eds): Leucocyte Typing III, White Cell Differentiation Antigens. Oxford University Press, Oxford. 1987; 1-1050. < NLM ID: 8913266 >
- Beverley PC: Human T cell subsets. Immunol Lett. 1987 Apr; 14(4):263-7. < PMID: 2953677 >
- Beverley PC, Merckenschlager M, Terry L: Phenotypic diversity of the CD45 antigen and its relationship to function. Immunol Suppl. 1988; 1:3-5. < PMID: 2968306 >
- Akbar AN, Terry L, Timms A, Beverley PC, Janossy G: Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T cells. J Immunol. 1988 Apr 1; 140(7):2171-8. < PMID: 2965180 >
- Terry LA, Brown MH, Beverley PC: The monoclonal antibody, UCHL1, recognizes a 180,000 MW component of the human leucocyte-common antigen, CD45. Immunology. 1988 Jun; 64(2):331-6. < PMID: 2455685 >
- Merckenschlager M, Terry L, Edwards R, Beverley PC: Limiting dilution analysis of proliferative responses in human lymphocyte populations defined by the monoclonal antibody UCHL1: implications for differential CD45 expression in T cell memory formation. Eur J Immunol. 1988 Nov; 18(11):1653-61. < PMID: 2974420 >
- Knapp W, Dorken B, Gilks W, Rieber EP, Schmidt RE, Stein H, von dem Borne AEGK (Eds): Leucocyte Typing IV. Oxford University Press, Oxford. 1989; 1-1820. < NLM ID: 8914679 >
- Doussis IA, Gatter KC, Mason DY: CD68 reactivity of non-macrophage derived tumours in cytological specimens. J Clin Pathol. 1993 Apr; 46(4):334-6. < PMID: 7684403 >
- Li FJ, Tsuyama N, Ishikawa H, Obata M, Abroun S, Liu S, Otsuyama K, Zheng X, Ma Z, Maki Y, Kawano MM: A rapid translocation of CD45RO but not CD45RA to lipid rafts in IL-6-induced proliferation in myeloma. Blood. 2005 Apr 15; 105(8):3295-302. < PMID: 15626731 >
- Dawes R, Petrova S, Liu Z, Wraith D, Beverley PC, Tchilian EZ: Combinations of CD45 isoforms are crucial for immune function and disease. J Immunol. 2006 Mar 15; 176(6):3417-25. < PMID: 16517710 >
- Cosenza-Nashat MA, Kim MO, Zhao ML, Suh HS, Lee SC: CD45 isoform expression in microglia and inflammatory cells in HIV-1 encephalitis.. Brain Pathol. 2006 Oct; 16(4):256-65. < PMID: 17107594 >
- Bijian K, Zhang L, Shen SH: Collagen-mediated survival signaling is modulated by CD45 in Jurkat T cells. Mol Immunol. 2007 Jul; 44(15):3682-90. < PMID: 17524482 >
- Desharnais P, Dupéré-Minier G, Hamelin C, Devine P, Bernier J: Involvement of CD45 in DNA fragmentation in apoptosis induced by mitochondrial perturbing agents. Apoptosis. 2008 Feb; 13(2):197-212. < PMID: 18157742 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
 Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com