

TECHNICAL DATA SHEET

CyFlow™ CD45RA Biotin Anti-Hu; Clone MEM-56

REF BC517380

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD45RA
Alternative Names	Leu18
Clone	MEM-56
Clonality	monoclonal
Format	Biotin
Host / Isotype	Mouse / IgG2b
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Human thymocytes and T lymphocytes

Specificity

The mouse monoclonal antibody MEM-56 recognizes CD45RA antigen, a 205-220 kDa single chain type I glycoprotein, isoform of the leukocyte common antigen (LCA). CD45RA is expressed on most of B lymphocytes, resting and native T lymphocytes, medullar thymocytes and monocytes.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for indirect immunofluorescence analysis by Flow Cytometry. Suggested working usage is 8.3 µg/ml. Indicated dilution is recommended starting point for use of this product, but working concentrations should be validated by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in Tris buffered saline (TBS) solution, pH ≈8.0, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD45RA is a high molecular weight isoform of a receptor-type protein tyrosine phosphatase, CD45 glycoprotein. CD45 is crucial in lymphocyte development and antigen signaling, serving as an important regulator of Src-family kinases, promotes cell survival by modulating integrin-mediated signal transduction pathway and is also involved in DNA fragmentation during apoptosis. CD45 isoforms differ in their extracellular domains, whereas they share identical transmembrane and cytoplasmic domains. These isoforms differ in their ability to translocate into the glycosphingolipid-enriched membrane domains and their expression depends on cell type and physiological state of the cell. CD45RA is expressed e.g. on naive T cells and normal plasma cells.

References

- Bazil V, Horejsi V, Baudys M, Kristofova H, Strominger JL, Kostka W, Hilgert I: Biochemical characterization of a soluble form of the 53-kDa monocyte surface antigen. Eur J Immunol. 1986 Dec; 16(12):1583-9. < PMID: 3493149 >
- Knapp W, Dorken B, Gilks W, Rieber EP, Schmidt RE, Stein H, von dem Borne AEGK (Eds): Leucocyte Typing IV. Oxford University Press, Oxford. 1989; 1-1820. < NLM ID: 8914679 >
- Bhunia AK, Arai T, Bulkley G, Chatterjee S: Lactosylceramide mediates tumor necrosis factor-alpha-induced intercellular adhesion molecule-1 (ICAM-1 expression and the adhesion of

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- neutrophil in human umbilical vein endothelial cells. J Biol Chem. 1998 Dec 18; 273(51):34349-57.
< PMID: 9852101 >
- Cortez-Gonzalez X, Sidney J, Adotevi O, Sette A, Millard F, Lemonnier F, Langlade-Demoyen P, Zanetti M: Immunogenic HLA-B7-restricted peptides of hTRT. Int Immunol. 2006 Dec; 18(12):1707-18.
< PMID: 17077179 >
 - Dawes R, Petrova S, Liu Z, Wraith D, Beverley PC, Tchilian EZ: Combinations of CD45 isoforms are crucial for immune function and disease. J Immunol. 2006 Mar 15; 176(6):3417-25.
< PMID: 16517710 >
 - Li C, Wong P, Pan T, Xiao F, Yin S, Chang B, Kang SC, Ironside J, Sy MS: Normal cellular prion protein is a ligand of selectins: binding requires Le(X) but is inhibited by sLe(X). Biochem J. 2007 Sep 1; 406(2):333-41. < PMID: 17497959 >
 - Desharnais P, Dupéré-Minier G, Hamelin C, Devine P, Bernier J: Involvement of CD45 in DNA fragmentation in apoptosis induced by mitochondrial perturbing agents. Apoptosis. 2008 Feb; 13(2):197-212. < PMID: 18157742 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com