

TECHNICAL DATA SHEET

CyFlow™ CD58 APC Anti-Hu; Clone MEM-63

REF AX001125

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD58
Alternative Names	LFA-3
Clone	MEM-63
Clonality	monoclonal
Format	APC
Host / Isotype	Mouse / IgG1
Species Reactivity	Human, Pig
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	NALM-6 human pre-B cell line

Specificity

The mouse monoclonal antibody MEM-63 recognizes CD58 antigen, a 40-70 kDa glycoprotein distributed over many tissues, leukocytes, erythrocytes, endothelial cells, epithelial cells and fibroblasts.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing Tris buffered saline (TBS) solution, pH ≈8.0, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD58 (LFA-3) is an immunoglobulin family adhesion molecule expressed by both hematopoietic and non-hematopoietic cells (often on antigen presenting cells) and serving as ligand of CD2. This interaction is important for T cell-mediated immunity. CD58 is expressed in transmembrane form and in GPI-anchored form; the later is constitutively associated with protein kinases whereas the transmembrane form activates kinase activity upon triggering. CD58 is a powerful tool for detection of minimal residual disease in acute lymphocytic leukemia, and for evaluation of liver damage related with hepatitis B.

References

- Kishimoto T, Goyert S, Kikutani H, Mason D, Miyasaka M, Moretta L, Ohno T, Okumura K, Shaw S, Springer TA, Sugamura K, Sugawara H, von dem Borne AEGK, Zola H (Eds): Leucocyte Typing VI. Garland Publishing Inc, New York. 1997; 1-1342. < NLM ID: 9712219 >
- Bayas MV, Schulten K, Leckband D: Forced detachment of the CD2-CD58 complex. Biophys J. 2003 Apr; 84(4):2223-33. < PMID: 12668431 >
- Veltroni M, De Zen L, Sanzari MC, Maglia O, Dworzak MN, Ratei R, Biondi A, Basso G, Gaipa G: I-BFM-ALL-FCM-MRD-Study Group: Expression of CD58 in normal, regenerating and leukemic bone marrow B cells: implications for the detection of minimal residual disease in acute lymphocytic leukemia. Haematologica. 2003 Nov; 88(11):1245-52. < PMID: 14607753 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Ariel O, Kukulansky T, Raz N, Hollander N: Distinct membrane localization and kinase association of the two isoforms of CD58. Cell Signal. 2004 Jun; 16(6):667-73. < PMID: 15093607 >
- Sheng L, Li J, Qi BT, Ji YQ, Meng ZJ, Xie M: Investigation on correlation between expression of CD58 molecule and severity of hepatitis B. World J Gastroenterol. 2006 Jul 14; 12(26):4237-40. < PMID: 16830383 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com