

TECHNICAL DATA SHEET

CyFlow™ CD64 Low Endotoxin Anti-Hu; Clone 10.1

REF AR810750

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD64
Alternative Names	FcγRI
Clone	10.1
Clonality	monoclonal
Format	Low Endotoxin
Host / Isotype	Mouse / IgG1
Species Reactivity	Human, Non-Human Primates
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Rheumatoid synovial fluid cells and fibronectin purified human monocytes

Specificity

The mouse monoclonal antibody 10.1 recognizes α subunit of CD64 antigen, a 72 kDa single chain type I glycoprotein, that is expressed on monocytes/macrophages, dendritic cells, and activated granulocytes.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Western blot
- Immunohistochemistry
- Immunocytochemistry
- Functional assays

Storage Buffer

The reagent is provided in azide-free phosphate buffered saline (PBS) solution, pH $\approx$ 7.4; 0.2 μ m filter sterilized. Endotoxin level is less than 0.01 EU/ μ g of the protein, as determined by the LAL test.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD64 (Fc γ RI) is a cell surface receptor for Fc region of IgG. It is composed of specific ligand binding α subunit and promiscuous γ subunit, which is indispensable for tyrosine-based signaling. However, even the α subunit can transduce signals leading to cellular effector functions. The isoform Fc γ RIa1 binds human IgG with high affinity, has limited myeloid cell distribution, and a relatively large intracellular domain. Products of related genes include Fc γ RIb and Fc γ RIc isoforms, but these specify low affinity IgG receptors if functionally expressed at all. Besides a role in antigen clearance, Fc γ RI (a1) can potentially enhance MHC class I and II antigen presentation in vitro and in vivo.

References

- Dougherty GJ, Selvendran Y, Murdoch S, Palmer DG, Hogg N: The human mononuclear phagocyte high-affinity Fc receptor, FcRI, defined by a monoclonal antibody, 10.1. Eur J Immunol. 1987 Oct; 17(10):1453-9. < PMID: 3500057 >
- Jayaram Y, Buckle AM, Hogg N: The Fc receptor, FcRI, and other activation molecules on human mononuclear phagocytes after treatment with interferon-gamma. Clin Exp Immunol. 1989 Mar; 75(3):414-20. < PMID: 2495203 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Hashimoto S, Yamada M, Motoyoshi K, Akagawa KS: Enhancement of macrophage colony-stimulating factor-induced growth and differentiation of human monocytes by interleukin-10. *Blood*. 1997 Jan 1; 89(1):315-21. < PMID: 8978307 >
- Fadlon E, Vordermeier S, Pearson TC, Mire-Sluis AR, Dumonde DC, Phillips J, Fishlock K, Brown KA: Blood polymorphonuclear leukocytes from the majority of sickle cell patients in the crisis phase of the disease show enhanced adhesion to vascular endothelium and increased expression of CD64. *Blood*. 1998 Jan 1; 91(1):266-74. < PMID: 9414294 >
- Sánchez-Torres C, García-Romo GS, Cornejo-Cortés MA, Rivas-Carvalho A, Sánchez-Schmitz G: CD16+ and CD16- human blood monocyte subsets differentiate in vitro to dendritic cells with different abilities to stimulate CD4+ T cells. *Int Immunol*. 2001 Dec; 13(12):1571-81. < PMID: 11717198 >
- Brichard B, Varis I, Latinne D, Deneys V, de Bruyere M, Leveugle P, Cornu G: Intracellular cytokine profile of cord and adult blood monocytes. *Bone Marrow Transplant*. 2001 May; 27(10):1081-6. < PMID: 11438825 >
- Beekman JM, Bakema JE, van der Linden J, Tops B, Hinten M, van Vugt M, van de Winkel JG, Leusen JH: Modulation of FcγRI (CD64) ligand binding by blocking peptides of periplakin. *J Biol Chem*. 2004 Aug 6; 279(32):33875-81. < PMID: 15161926 >
- Devaraj S, Du Clos TW, Jialal I: Binding and internalization of C-reactive protein by Fcγ receptors on human aortic endothelial cells mediates biological effects. *Arterioscler Thromb Vasc Biol*. 2005 Jul; 25(7):1359-63. < PMID: 15860734 >
- Roura-Mir C, Wang L, Cheng TY, Matsunaga I, Dascher CC, Peng SL, Fenton MJ, Kirschning C, Moody DB: Mycobacterium tuberculosis regulates CD1 antigen presentation pathways through TLR-2. *J Immunol*. 2005 Aug 1; 175(3):1758-66. < PMID: 16034117 >
- Devaraj S, Davis B, Simon SI, Jialal I: CRP promotes monocyte-endothelial cell adhesion via Fcγ receptors in human aortic endothelial cells under static and shear flow conditions. *Am J Physiol Heart Circ Physiol*. 2006 Sep; 291(3):H1170-6. < PMID: 16603696 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com