

TECHNICAL DATA SHEET

CyFlow™ CD79a FITC Anti-Hu/Ms/Rt; Clone HM57

REF CV675949

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD79a
Alternative Names	IgA, MB-1, MB1
Clone	HM57
Clonality	monoclonal
Format	FITC
Host / Isotype	Mouse / IgG1
Species Reactivity	Human Mouse Rat, Pig Cow Horse Rabbit Guinea pig Opossum Chicken
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Synthetic peptide corresponding to amino acids 202-216 of human CD79a

Specificity

The mouse monoclonal antibody HM57 recognizes CD79a, a 40-45 kDa subunit of B cell antigen-specific receptor (BCR) and its early developmental forms.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD79a (Ig α, MB1) forms disulfide-linked heterodimer with CD79b (Ig β, B29). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immunoglobulin they constitute B-cell antigen-specific receptor (BCR). CD79a and b are the first components of BCR that are expressed developmentally. They appear on pro-B cells in association with the endoplasmic reticulum chaperone calnexin. Subsequently, in pre-B cells, CD79 heterodimer is associated with λ5-VpreB surrogate immunoglobulin and later with antigen-specific surface immunoglobulins. At the plasma cell stage, CD79a is present as an intracellular component. CD79a/b complex interacts with Src-family tyrosine kinase Lyn, which phosphorylates its cytoplasmic ITAM motives to form docking sites for downstream signaling.

References

- Mason DY, Cordell JL, Tse AG, van Dongen JJ, van Noesel CJ, Micklem K, Pulford KA, Valensi F, Comans-Bitter WM, Borst J, et al.: The IgM-associated protein mb-1 as a marker of normal and neoplastic B cells. J Immunol. 1991 Dec 1; 147(11):2474-82. < PMID: 1747162 >
- van Noesel CJ, van Lier RA, Cordell JL, Tse AG, van Schijndel GM, de Vries EF, Mason DY, Borst J: The membrane IgM-associated heterodimer on human B cells is a newly defined B cell antigen that contains the protein product of the mb-1 gene. J Immunol. 1991 Jun 1; 146(11):3881-8. < PMID: 2033258 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Mason DY, van Noesel CJ, Cordell JL, Comans-Bitter WM, Micklem K, Tse AG, van Lier RA, van Dongen JJ: The B29 and mb-1 polypeptides are differentially expressed during human B cell differentiation. *Eur J Immunol.* 1992 Oct; 22(10):2753-6. < PMID: 1396979 >
- Doussis IA, Gatter KC, Mason DY: CD68 reactivity of non-macrophage derived tumours in cytological specimens. *J Clin Pathol.* 1993 Apr; 46(4):334-6. < PMID: 7684403 >
- Jones M, Cordell JL, Beyers AD, Tse AG, Mason DY: Detection of T and B cells in many animal species using cross-reactive anti-peptide antibodies. *J Immunol.* 1993 Jun 15; 150(12):5429-35. < PMID: 8515069 >
- Mason DY, Cordell JL, Brown MH, Borst J, Jones M, Pulford K, Jaffe E, Ralfkiaer E, Dallenbach F, Stein H, et al.: CD79a: a novel marker for B-cell neoplasms in routinely processed tissue samples. *Blood.* 1995 Aug 15; 86(4):1453-9. < PMID: 7632952 >
- Kishimoto T, Goyert S, Kikutani H, Mason D, Miyasaka M, Moretta L, Ohno T, Okumura K, Shaw S, Springer TA, Sugamura K, Sugawara H, von dem Borne AEGK, Zola H (Eds): *Leucocyte Typing VI*. Garland Publishing Inc, New York. 1997; 1-1342. < NLM ID: 9712219 >
- Bannish G, Fuentes-Pananá EM, Cambier JC, Pear WS, Monroe JG: Ligand-independent signaling functions for the B lymphocyte antigen receptor and their role in positive selection during B lymphopoiesis. *J Exp Med.* 2001 Dec 3; 194(11):1583-96. < PMID: 11733573 >
- Pike KA, Iacampo S, Friedmann JE, Ratcliffe MJ: The cytoplasmic domain of Ig alpha is necessary and sufficient to support efficient early B cell development. *J Immunol.* 2004 Feb 15; 172(4):2210-8. < PMID: 14764688 >
- Fuentes-Pananá EM, Bannish G, Shah N, Monroe JG: Basal Igalpha/Igbeta signals trigger the coordinated initiation of pre-B cell antigen receptor-dependent processes. *J Immunol.* 2004 Jul 15; 173(2):1000-11. < PMID: 15240688 >
- Fuentes-Pananá EM, Bannish G, van der Voort D, King LB, Monroe JG: Ig alpha/Ig beta complexes generate signals for B cell development independent of selective plasma membrane compartmentalization. *J Immunol.* 2005 Feb 1; 174(3):1245-52. < PMID: 15661879 >
- Fuentes-Pananá EM, Bannish G, Karnell FG, Trembl JF, Monroe JG: Analysis of the individual contributions of Igalpha (CD79a)- and Igbeta (CD79b)-mediated tonic signaling for bone marrow B cell development and peripheral B cell maturation. *J Immunol.* 2006 Dec 1; 177(11):7913-22. < PMID: 17114463 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Faldyna M, Samankova P, Leva L, Cerny J, Oujezdska J, Rehakova Z, Sinkora J: Cross-reactive anti-human monoclonal antibodies as a tool for B-cell identification in dogs and pigs. Vet Immunol Immunopathol. 2007 Sep 15; 119(1-2):56-62. < PMID: 17673300 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com