

TECHNICAL DATA SHEET

CyFlow™ CD79a PE Anti-Hu/Ms/Rt; Clone HM47

REF CU479884

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD79a
Alternative Names	IgA, MB-1, MB1
Clone	HM47
Clonality	monoclonal
Format	PE
Host / Isotype	Mouse / IgG1
Species Reactivity	Human Mouse Rat, Non-Human Primates Pig Cow Horse Dog Rabbit Guinea pig Chicken
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Synthetic peptide corresponding to C terminal amino acids 208-222 of human CD79a

Specificity

The mouse monoclonal antibody HM57 recognizes CD79a, a 40-45 kDa subunit of B cell antigen-specific receptor (BCR) and its early developmental forms.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD79a (Ig α, MB1) forms disulfide-linked heterodimer with CD79b (Ig β, B29). They both are transmembrane proteins with extended cytoplasmic domains containing immunoreceptor tyrosine activation motives (ITAMs), and together with cell surface immunoglobulin they constitute B-cell antigen-specific receptor (BCR). CD79a and b are the first components of BCR that are expressed developmentally. They appear on pro-B cells in association with the endoplasmic reticulum chaperone calnexin. Subsequently, in pre-B cells, CD79 heterodimer is associated with λ5-VpreB surrogate immunoglobulin and later with antigen-specific surface immunoglobulins. At the plasma cell stage, CD79a is present as an intracellular component. CD79a/b complex interacts with Src-family tyrosine kinase Lyn, which phosphorylates its cytoplasmic ITAM motives to form docking sites for downstream signaling.

References

- Mason DY, Cordell JL, Tse AG, van Dongen JJ, van Noesel CJ, Micklem K, Pulford KA, Valensi F, Comans-Bitter WM, Borst J, et al.: The IgM-associated protein mb-1 as a marker of normal and neoplastic B cells. J Immunol. 1991 Dec 1; 147(11):2474-82. < PMID: 1747162 >
- Rassenti LZ, Kipps TJ: Expression of Ig-beta (CD79b) by chronic lymphocytic leukemia B cells that lack immunoglobulin heavy-chain allelic exclusion. Blood. 2000 Apr 15; 95(8):2725-7. < PMID: 10753858 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Torlakovic E, Torlakovic G: B-cell markers in lymphocyte predominance Hodgkin disease. Arch Pathol Lab Med. 2002 Jul; 126(7):862-3. < PMID: 12123231 >
- Islas-Ohlmayer M, Padgett-Thomas A, Domiati-Saad R, Melkus MW, Cravens PD, Martin Mdel P, Netto G, Garcia JV: Experimental infection of NOD/SCID mice reconstituted with human CD34+ cells with Epstein-Barr virus. J Virol. 2004 Dec; 78(24):13891-900. < PMID: 15564497 >
- Fernandez NJ, West KH, Jackson ML, Kidney BA: Immunohistochemical and histochemical stains for differentiating canine cutaneous round cell tumors.. Vet Pathol. 2005 Jul; 42(4):437-45. < PMID: 16006603 >
- Bhargava P, Kallakury BV, Ross JS, Azumi N, Bagg A: CD79a is heterogeneously expressed in neoplastic and normal myeloid precursors and megakaryocytes in an antibody clone-dependent manner. Am J Clin Pathol. 2007 Aug; 128(2):306-13. < PMID: 17638667 >
- Zhao XF, Hassan A, Perry A, Ning Y, Stass SA, Dehner LP: C-MYC rearrangements are frequent in aggressive mature B-Cell lymphoma with atypical morphology. Int J Clin Exp Pathol. 2008 Jan 1; 1(1):65-74. < PMID: 18784824 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com