

TECHNICAL DATA SHEET

CyFlow™ CD86 APC Anti-Hu; Clone BU63

REF AX099363

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD86
Alternative Names	B70, B7-2
Clone	BU63
Clonality	monoclonal
Format	APC
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	B-lymphoblastoid cell line ARH 77

Specificity

The mouse monoclonal antibody BU63 recognizes CD86 antigen, a 70 kDa type I transmembrane glycoprotein of immunoglobulin supergene family, expressed on professional antigen-presenting cells, such as dendritic cells, macrophages or activated B lymphocytes.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD86 (B7-2) and CD80 (B7-1) are ligands of T cell critical costimulatory molecule CD28 and of an inhibitory receptor CD152 (CTLA-4). Both B7 molecules are expressed on professional antigen-presenting cells and are essential for T cell activation, both molecules can also substitute for each other in this process. The question what are the differences in CD80 and CD86 competency has not been fully elucidated yet; there are still conflicts in results about their respective roles in initiation or sustaining of the T cell immune response.

References

- Doussis IA, Gatter KC, Mason DY: CD68 reactivity of non-macrophage derived tumours in cytological specimens. J Clin Pathol. 1993 Apr; 46(4):334-6. < PMID: 7684403 >
- Engel P, Gribben JG, Freeman GJ, Zhou LJ, Nozawa Y, Abe M, Nadler LM, Wakasa H, Tedder TF: The B7-2 (B70) costimulatory molecule expressed by monocytes and activated B lymphocytes is the CD86 differentiation antigen. Blood. 1994 Sep 1; 84(5):1402-7. < PMID: 7520767 >
- Caux C, Vanbervliet B, Massacrier C, Azuma M, Okumura K, Lanier LL, Banchereau J: B70/B7-2 is identical to CD86 and is the major functional ligand for CD28 expressed on human dendritic cells. J Exp Med. 1994 Nov 1; 180(5):1841-7. < PMID: 7525840 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Mauri D, Wyss-Coray T, Gallati H, Pichler WJ: Antigen-presenting T cells induce the development of cytotoxic CD4+ T cells (I): Involvement of the CD80-CD28 adhesion molecules. J Immunol. 1995 Jul 1; 155(1):118-27. < PMID: 7541409 >
- Kishimoto T, Goyert S, Kikutani H, Mason D, Miyasaka M, Moretta L, Ohno T, Okumura K, Shaw S, Springer TA, Sugamura K, Sugawara H, von dem Borne AEGK, Zola H (Eds): Leucocyte Typing VI. Garland Publishing Inc, New York. 1997; 1-1342. < NLM ID: 9712219 >
- Vasilevko V, Ghochikyan A, Holterman MJ, Agadjanyan MG: CD80 (B7-1 and CD86 (B7-2 are functionally equivalent in the initiation and maintenance of CD4+ T-cell proliferation after activation with suboptimal doses of PHA. DNA Cell Biol. 2002 Mar; 21(3):137-49. < PMID: 12015893 >
- Zhan H, Towler HM, Calder VL: The immunomodulatory role of human conjunctival epithelial cells. Invest Ophthalmol Vis Sci. 2003 Sep; 44(9):3906-10. < PMID: 12939308 >
- Giguere JF, Bounou S, Paquette JS, Madrenas J, Tremblay MJ: Insertion of host-derived costimulatory molecules CD80 (B7.1) and CD86 (B7.2) into human immunodeficiency virus type 1 affects the virus life cycle. J Virol. 2004 Jun; 78(12):6222-32. < PMID: 15163715 >
- Yadav D, Judkowski V, Flodstrom-Tullberg M, Sterling L, Redmond WL, Sherman L, Sarvetnick N: B7-2 (CD86 controls the priming of autoreactive CD4 T cell response against pancreatic islets. J Immunol. 2004 Sep 15; 173(6):3631-9. < PMID: 15356107 >
- Chan A, Baird M, Mercer AA, Fleming SB: Maturation and function of human dendritic cells are inhibited by orf virus-encoded interleukin-10. J Gen Virol. 2006 Nov; 87(11):3177-81. < PMID: 17030850 >
- Kolar GR, Mehta D, Pelayo R, Capra JD: A novel human B cell subpopulation representing the initial germinal center population to express AID. Blood. 2007 Mar 15; 109(6):2545-52. < PMID: 17132718 >
- Thomas IJ, Petrich de Marquesini LG, Ravanani R, Smith RM, Guerder S, Flavell RA, Wraith DC, Wen L, Wong FS: CD86 has sustained costimulatory effects on CD8 T cells. J Immunol. 2007 Nov 1; 179(9):5936-46. < PMID: 17947667 >
- Eri R, Kodumudi KN, Summerlin DJ, Srinivasan M: Suppression of colon inflammation by CD80 blockade: evaluation of two murine models of inflammatory bowel disease. Inflamm Bowel Dis. 2008 Jan 9; < PMID: 21061388 >
- Hovden AO, Karlsen M, Jonsson R, Aarstad HJ, Appel S: Maturation of monocyte derived dendritic cells with OK432 boosts IL-12p70 secretion and conveys strong T-cell responses. BMC Immunol. 2011 Jan 5; 12:2. < PMID: 21208424 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com