

TECHNICAL DATA SHEET

CyFlow™ CD8a Purified Anti-Ms; Clone 53-6.7

REF CC027137

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD8a
Alternative Names	T8, Leu2
Clone	53-6.7
Clonality	monoclonal
Format	Purified
Host / Isotype	Rat / IgG2a
Species Reactivity	Mouse
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Mouse spleen cells

Specificity

The rat monoclonal antibody 53-6.7 recognizes mouse CD8a antigen, a 32-34 kDa α chain of the CD8 antigen.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry
- Immunoprecipitation
- Immunohistochemistry
- Immunocytochemistry
- Functional assays

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

The CD8a (CD8 α) subunit of CD8 T cell coreceptor is expressed in CD8 α/β heterodimers on majority of MHC I-restricted conventional T cells and thymocytes and in CD8 α/α homodimers on subsets of memory T cells, intraepithelial lymphocytes, NK cells, macrophages and dendritic cells. Regulation of CD8 beta level on T cell surface seems to be an important mechanism to control their effector function. Assembly of CD8 α/β but not α/α dimers is connected with formation or localization to the lipid rafts. Recruiting triggered TCR complexes to these membrane microdomains as well as affinity of TCR to MHC I is modulated by CD8, thereby affecting the functional diversity of the TCR signaling.

References

- Ledbetter JA, Herzenberg LA: Xenogeneic monoclonal antibodies to mouse lymphoid differentiation antigens. Immunol Rev. 1979; 47:63-90. < PMID: 398327 >
- Ledbetter JA, Rouse RV, Micklem HS, Herzenberg LA: T cell subsets defined by expression of Lyt-1,2,3 and Thy-1 antigens: Two-parameter immunofluorescence and cytotoxicity analysis with monoclonal antibodies modifies current views. J Exp Med. 1980 Aug 1; 152(2):280-95. < PMID: 6156984 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Takahashi K, Nakata M, Tanaka T, Adachi H, Nakauchi H, Yagita H, Okumura K: CD4 and CD8 regulate interleukin 2 responses of T cells. Proc Natl Acad Sci USA. 1992 Jun 15; 89(12):5557-61. < PMID: 1608966 >
- Grabbe S, Varga G, Beissert S, Steinert M, Pendl G, Seeliger S, Bloch W, Peters T, Schwarz T, Sunderkötter C, Scharffetter-Kochanek K: Beta2 integrins are required for skin homing of primed T cells but not for priming naïve T cells. J Clin Invest. 2002 Jan; 109(2):183-92. < PMID: 11805130 >
- Hata H, Sakaguchi N, Yoshitomi H, Iwakura Y, Sekikawa K, Azuma Y, Kanai C, Moriizumi E, Nomura T, Nakamura T, Sakaguchi S: Distinct contribution of IL-6, TNF-alpha, IL-1, and IL-10 to T cell-mediated spontaneous autoimmune arthritis in mice. J Clin Invest. 2004 Aug; 114(4):582-8. < PMID: 15314695 >
- Ko SY, Ko HJ, Chang WS, Park SH, Kweon MN, Kang CY: alpha-Galactosylceramide can act as a nasal vaccine adjuvant inducing protective immune responses against viral infection and tumor. J Immunol. 2005 Sep 1; 175(5):3309-17. < PMID: 16116223 >
- Bouwer HG, Alberti-Segui C, Montfort MJ, Berkowitz ND, Higgins DE: Directed antigen delivery as a vaccine strategy for an intracellular bacterial pathogen. Proc Natl Acad Sci USA. 2006 Mar 28; 103(13):5102-7. < PMID: 16549792 >
- Kamimura D, Sawa Y, Sato M, Agung E, Hirano T, Murakami M: IL-2 in vivo activities and antitumor efficacy enhanced by an anti-IL-2 mAb. J Immunol. 2006 Jul 1; 177(1):306-14. < PMID: 16785526 >
- Mochimaru H, Usui T, Yaguchi T, Nagahama Y, Hasegawa G, Usui Y, Shimmura S, Tsubota K, Amano S, Kawakami Y, Ishida S: Suppression of alkali burn-induced corneal neovascularization by dendritic cell vaccination targeting VEGF receptor 2. Invest Ophthalmol Vis Sci. 2008 May; 49(5):2172-7. < PMID: 18263815 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com