

TECHNICAL DATA SHEET

CyFlow™ CD8b PE Anti-Rt; Clone 341

REF AJ377672

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD8b
Alternative Names	—
Clone	341
Clonality	monoclonal
Format	PE
Host / Isotype	Mouse / IgG1
Species Reactivity	Rat
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [0.5 mg/ml]
Immunogen	CD8 positive Wistar rat splenic T cell hybridomas

Specificity

The mouse monoclonal antibody 341 (also known as 34.1) recognizes rat CD8b antigen, a 32-34 kDa β chain of the CD8 coreceptor, expressed on T cell subsets and some other cell types, such as macrophages.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Working concentrations should be determined by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

The CD8b (CD8 β) subunit of CD8 T cell coreceptor is expressed in CD8 α/β heterodimers on majority of MHC I-restricted conventional T cells and thymocytes and in CD8 α/α homodimers on subsets of memory T cells, intraepithelial lymphocytes, NK cells, macrophages, mast cells, and dendritic cells. Regulation of CD8 beta level on T cell surface seems to be an important mechanism to control their effector function. Assembly of CD8 α/β but not α/α dimers is connected with formation or localization to the lipid rafts. Recruiting triggered TCR complexes to these membrane microdomains as well as affinity of TCR to MHC I is modulated by CD8, thereby affecting the functional diversity of the TCR signaling.

References

- Torres-Nagel N, Kraus E, Brown MH, Tiefenthaler G, Mitnacht R, Williams AF, Hünig T: Differential thymus dependence of rat CD8 isoform expression. Eur J Immunol. 1992 Nov; 22(11):2841-8. < PMID: 1358623 >
- Kühnlein P, Park JH, Herrmann T, Elbe A, Hünig T: Identification and characterization of rat gamma/delta T lymphocytes in peripheral lymphoid organs, small intestine, and skin with a monoclonal antibody to a constant determinant of the gamma/delta T cell receptor. J Immunol. 1994 Aug1; 153(3):979-86. < PMID: 8027567 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Kraus E, Lambracht D, Wonigeit K, Hünig T: Negative regulation of rat natural killer cell activity by major histocompatibility complex class I recognition. Eur J Immunol. 1996 Nov; 26(11):2582-6. < PMID: 8921942 >
- Hirji N, Lin TJ, Befus AD: A novel CD8 molecule expressed by alveolar and peritoneal macrophages stimulates nitric oxide production. J Immunol. 1997 Feb 15; 158(4):1833-40. < PMID: 9029123 >
- Lin TJ, Hirji N, Nohara O, Stenton GR, Gilchrist M, Befus AD: Mast cells express novel CD8 molecules that selectively modulate mediator secretion. J Immunol. 1998 Dec 1; 161(11):6265-72. < PMID: 9834115 >
- Hirji N, Lin TJ, Bissonnette E, Belosevic M, Befus AD: Mechanisms of macrophage stimulation through CD8: macrophage CD8alpha and CD8beta induce nitric oxide production and associated killing of the parasite Leishmania major. J Immunol. 1998 Jun 15; 160(12):6004-11. < PMID: 9637515 >
- Ellerman KE, Like AA: Islet cell membrane antigens activate diabetogenic CD4+ T-cells in the BB/Wor rat. Diabetes. 1999 May; 48(5):975-82. < PMID: 10331400 >
- Nohara O, Kulka M, Déry RE, Wills FL, Hirji NS, Gilchrist M, Befus AD: Regulation of CD8 expression in mast cells by exogenous or endogenous nitric oxide. J Immunol. 2001 Nov 15; 167(10):5935-9. < PMID: 11698471 >
- Mabarrack NH, Turner NL, Mayrhofer G: Recent thymic origin, differentiation, and turnover of regulatory T cells. J Leukoc Biol. 2008 Nov; 84(5):1287-97. < PMID: 18682578 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com