

TECHNICAL DATA SHEET

CyFlow™ CD98 FITC Anti-Hu; Clone MEM-108

REF BV547235

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD98
Alternative Names	4F2, FRP-1
Clone	MEM-108
Clonality	monoclonal
Format	FITC
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Raji cells: human Burkitt's lymphoma cell line

Specificity

The mouse monoclonal antibody MEM-108 recognizes CD98 antigen, a 125 kDa disulfide-linked heterodimer (80 kDa glycosylated heavy chain + 45 kDa non-glycosylated light chain). CD98 is expressed

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

on T lymphocytes (upon activation) and activated NK cells; it is also present at low levels on B lymphocytes, NK cells, monocytes and platelets.

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD98 (4F2) is a type II transmembrane glycoprotein which serves as the heavy chain of the heterodimeric amino acid transporters (HATs). CD98, linked to various light chains by disulfide bond, is responsible for cell surface expression and basolateral localization of this transporter complex in polarized epithelial cells and also interacts with β1 integrins and increases their affinity for ligand. Besides its roles in amino acid transport, CD98 is thus involved in cell fusion and activation. It is implicated in regulation of cellular differentiation, growth and apoptosis.

References

- Kishimoto T, Goyert S, Kikutani H, Mason D, Miyasaka M, Moretta L, Ohno T, Okumura K, Shaw S, Springer TA, Sugamura K, Sugawara H, von dem Borne AEGK, Zola H (Eds): Leucocyte Typing VI. Garland Publishing Inc, New York. 1997; 1-1342. < NLM ID: 9712219 >
- Cho JY, Skubitz KM, Katz DR, Chain BM: CD98-dependent homotypic aggregation is associated with translocation of protein kinase Cdelta and activation of mitogen-activated protein kinases. Exp Cell Res. 2003 May 15; 286(1):1-11. < PMID: 12729789 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Liu X, Charrier L, Gewirtz A, Sitaraman S, Merlin D: CD98 and intracellular adhesion molecule I regulate the activity of amino acid transporter LAT-2 in polarized intestinal epithelia. J Biol Chem. 2003 Jun 27; 278(26):23672-7. < PMID: 12716892 >
- Cai S, Bulus N, Fonseca-Siesser PM, Chen D, Hanks SK, Pozzi A, Zent R: CD98 modulates integrin beta1 function in polarized epithelial cells. J Cell Sci. 2005 Mar 1; 118(5):889-99. < PMID: 15713750 >
- Dalton P, Christian HC, Redman CW, Sargent IL, Boyd CA: Differential effect of cross-linking the CD98 heavy chain on fusion and amino acid transport in the human placental trophoblast (BeWo) cell line. Biochim Biophys Acta. 2007 Mar; 1768(3):401-10. < PMID: 17258169 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com