

TECHNICAL DATA SHEET

CyFlow™ CD99 APC **Anti-Hu; Clone 3B2/TA8**

REF BM091620

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	CD99
Alternative Names	MIC2, E2
Clone	3B2/TA8
Clonality	monoclonal
Format	APC
Host / Isotype	Mouse / IgG2a
Species Reactivity	Human
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Human thymocytes

Specificity

The mouse monoclonal antibody 3B2/TA8 recognizes CD99 antigen, a 32 kDa sialoglycoprotein expressed on many cell types, with particularly strong expression on Ewing's sarcoma and peripheral

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

primitive neuroectodermal tumors. Within the hematopoietic system, CD99 is expressed on virtually all cell types except granulocytes.

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

CD99 is a ubiquitous transmembrane type I sialoglycoprotein of a unique and poorly characterized protein family. CD99 is heavily O-glycosylated and was described as a T cell costimulator and strong activator of integrin-mediated actin cytoskeleton assembly, promoting cell adhesion and homotypic aggregation, immediate arrest on an inflamed vascular endothelium, and cell migration through it. Ligation of CD99 under some conditions can lead to apoptosis. Originally CD99 was described as a human thymus leukemia antigen, an Ewing's sarcoma-specific membrane marker, and an adhesion molecule involved in spontaneous rosette formation of T cells with erythrocytes.

References

- Waclavicek M, Majdic O, Stulnig T, Berger M, Sunder-Plassmann R, Zlabinger GJ, Baumruker T, Stöckl J, Ebner C, Knapp W, Pickl WF: CD99 engagement on human peripheral blood T cells results in TCR/CD3-dependent cellular activation and allows for Th1-restricted cytokine production. J Immunol. 1998 Nov 1; 161(9):4671-8. < PMID: 9794396 >
- Pickl WF, Pimentel-Muiños FX, Seed B: Lipid rafts and pseudotyping. J Virol. 2001 Aug; 75(15):7175-83. < PMID: 11435598 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Kish DD, Gorbachev AV, Fairchild RL: CD8+ T cells produce IL-2, which is required for CD(4+)CD25+ T cell regulation of effector CD8+ T cell development for contact hypersensitivity responses. J Leukoc Biol. 2005 Sep; 78(3):725-35. < PMID: 16000396 >
- Kueng HJ, Leb VM, Haiderer D, Raposo G, Thery C, Derdak SV, Schmetterer KG, Neunkirchner A, Sillaber C, Seed B, Pickl WF: General strategy for decoration of enveloped viruses with functionally active lipid-modified cytokines. J Virol. 2007 Aug; 81(16):8666-76. < PMID: 17537846 >
- Brémond A, Meynet O, Mahiddine K, Coito S, Tichet M, Scotlandi K, Breitmayer JP, Gounon P, Gleeson PA, Bernard A, Bernard G: Regulation of HLA class I surface expression requires CD99 and p230/golgin-245 interaction. Blood. 2009 Jan 8; 113(2):347-57. < PMID: 18849489 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com