

TECHNICAL DATA SHEET

CyFlow™ CK-18 FITC Clone C-04

REF BW627672

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	Cytokeratin 18
Alternative Names	—
Clone	C-04
Clonality	monoclonal
Format	FITC
Host / Isotype	Mouse / IgG1
Species Reactivity	Mammalian
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [0.1 mg/ml]
Immunogen	Cytoskeleton preparation of epidermal carcinoma cell line A431

Specificity

The mouse monoclonal antibody C-04 recognizes Cytokeratin 18, a 45 kDa member of intermediate filaments subfamily represented in epithelial tissues.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Working concentrations should be determined by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

Cytokeratins are a subfamily of intermediate filaments and characterized by remarkable biochemical diversity. Cytokeratins are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The individual cytokeratin polypeptides are designated 1 to 20 and divided into the type I (acidic cytokeratins 9-20) and type II (basic to neutral cytokeratins 1-8) families. Cytokeratin 18 belongs to type I family (acidic cytokeratins).

References

- Kovarik J, Rejthar A, Lauerova L, Vojtesek B, Bartkova J: Monoclonal antibodies against individual cytokeratins in the detection of metastatic spread. Int J Cancer Suppl. 1988; 3:50-5. < PMID: 2463228 >
- Lauerova L, Kovarik J, Bartek J, Rejthar A, Vojtesek B: Novel monoclonal antibodies defining epitope of human cytokeratin 18 molecule. Hybridoma. 1988 Oct; 7(5):495-504. < PMID: 2461901 >
- Vojtěšek B, Stasková Z, Nenutil R, Lauerová L, Kovarik J, Rejthar A, Bártek J, Bártek J.: Monoclonal antibodies recognizing different epitopes of cytokeratin No.18. Folia Biol (Praha). 1989; 35(6):373-82. < PMID: 2483834 >
- Taylor-Papadimitriou J, Stampfer M, Bartek J, Lewis A, Boshell M, Lane EB, Leigh IM: Keratin expression in human mammary epithelial cells cultured from normal and malignant tissue: relation to in vivo phenotypes and influence of medium. J Cell Sci. 1989 Nov; 94 (3):403-13. < PMID: 2483723 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Kovarik J, et al.: J Tumor Marker Oncol. 1990; 5:219. < No PMID >
- Bartek J, Vojtesek B, Staskova Z, Bartkova J, Kerekes Z, Rejthar A, Kovarik J: A series of 14 new monoclonal antibodies to keratins: characterization and value in diagnostic histopathology. J Pathol. 1991 Jul; 164(3):215-24. < PMID: 1716305 >
- Cap J, Cerman J, Nemecek S, Marekova M, Hana V, Frysak Z: The influence of treatment with somatostatin analogues on morphology, proliferative and apoptotic activity in GH-secreting pituitary adenomas. J Clin Neurosci. 2003 Jul; 10(4):444-8. < PMID: 12852883 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com