

TECHNICAL DATA SHEET

CyFlow™ CK-Pan PE Clone C-11

REF AJ818165

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	Cytokeratin (Pan-reactive)
Alternative Names	—
Clone	C-11
Clonality	monoclonal
Format	PE
Host / Isotype	Mouse / IgG1
Species Reactivity	Mammalian
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [0.1 mg/ml]
Immunogen	Keratin-enriched preparation from human epidermoid carcinoma cell line A431

Specificity

The mouse monoclonal antibody C-11 recognizes Cytokeratin peptides 4, 5, 6, 8, 10, 13, 18. Cytokeratins are a member of intermediate filaments subfamily represented in epithelial tissues.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Working concentrations should be determined by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

Cytokeratins are a subfamily of intermediate filaments and characterized by remarkable biochemical diversity. Cytokeratins are represented in epithelial tissues by at least 20 different polypeptides, molecular weight between 40 kDa and 68 kDa. The individual cytokeratin polypeptides are designated 1 to 20 and divided into the type I (acidic cytokeratins 9-20) and type II (basic to neutral cytokeratins 1-8) families.

References

- Kovarik J, Rejthar A, Lauerova L, Vojtesek B, Bartkova J: Monoclonal antibodies against individual cytokeratins in the detection of metastatic spread. Int J Cancer Suppl. 1988; 3:50-5. < PMID: 2463228 >
- Vojtěšek B, Stasková Z, Nenutil R, Lauerová L, Kovarik J, Rejthar A, Bártková J, Bártek J.: Monoclonal antibodies recognizing different epitopes of cytokeratin No.18. Folia Biol (Praha). 1989; 35(6):373-82. < PMID: 2483834 >
- Bartek J, Vojtesek B, Staskova Z, Bartkova J, Kerekes Z, Rejthar A, Kovarik J: A series of 14 new monoclonal antibodies to keratins: characterization and value in diagnostic histopathology. J Pathol. 1991 Jul; 164(3):215-24. < PMID: 1716305 >
- Hamakawa H, Sumida T, Tanioka H, Sogawa K, Yamada T: Extraction of cytokeratin from the human submandibular gland and its electrophoretic analysis. Res Commun Mol Pathol Pharmacol. 1998 Aug; 101(2):115-26. < PMID: 9821208 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Broekema M, Harmsen MC, Koerts JA, Petersen AH, van Luyn MJ, Navis G, Popa ER: Determinants of tubular bone marrow-derived cell engraftment after renal ischemia/reperfusion in rats. Kidney Int. 2005 Dec; 68(6):2572-81. < PMID: 16316332 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com