

TECHNICAL DATA SHEET

CyFlow™ Helios Purified Anti-Hu/Ms; Clone 22F6

REF CQ790210

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	Helios
Alternative Names	IKZF2
Clone	22F6
Clonality	monoclonal
Format	Purified
Host / Isotype	Hamster / IgG
Species Reactivity	Human Mouse
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	Peptide corresponding to the amino acids 51-107 of Helios

Specificity

The armenian hamster monoclonal antibody 22F6 recognizes Helios antigen, a transcription factor expressed in some hematopoietic stem cells, and at high levels in thymic-derived regulatory T cells.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

Based on published sources, this antibody is suitable for the following applications:

- Flow cytometry

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

Helios (IKZF2; Ikaros family zinc finger protein 2) is a hematopoietic-specific transcription factor involved in the regulation of lymphocyte development, together with other members of this family, such as Aiolos and Ikaros. Helios forms homo- and heterodimers with these proteins and is thought to function predominantly in early hematopoietic development. Expression of Helios, Aiolos and Ikaros is restricted to cells of the hematopoietic system, whereas other family members, Eos and Pegassus, are more widely expressed. Helios is expressed at early stages of thymocyte development. In mature T cells, Helios has been strongly associated with Treg cells.

References

- Thornton AM, Korty PE, Tran DQ, Wohlfert EA, Murray PE, Belkaid Y, Shevach EM: Expression of Helios, an Ikaros transcription factor family member, differentiates thymic-derived from peripherally induced Foxp3+ T regulatory cells. J Immunol. 2010 Apr 1; 184(7):3433-41. < PMID: 20181882 >
- Pinheiro D, Singh Y, Grant CR, Appleton RC, Sacchini F, Walker KR, Chadbourne AH, Palmer CA, Armitage-Chan E, Thompson I, Williamson L, Cunningham F, Garden OA: Phenotypic and functional characterization of a CD4(+) CD25(high) FOXP3(high) regulatory T-cell population in the dog. Immunology. 2011 Jan; 132(1):111-22. < PMID: 20880379 >
- Serre K, Bénézech C, Desanti G, Bobat S, Toellner KM, Bird R, Chan S, Kastner P, Cunningham AF, MacLennan IC, Mohr E: Helios is associated with CD4 T cells differentiating to T helper 2 and follicular helper T cells in vivo independently of Foxp3 expression. PLoS One. 2011; 6(6):e20731. < PMID: 25877073 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Daley SR, Hu DY, Goodnow CC: Helios marks strongly autoreactive CD4+ T cells in two major waves of thymic deletion distinguished by induction of PD-1 or NF- κ B. J Exp Med. 2013 Feb 11; 210(2):269-85. < PMID: 23337809 >
- McIver Z, Melenhorst JJ, Wu C, Grim A, Ito S, Cho I, Hensel N, Battiwalla M, Barrett AJ: Donor lymphocyte count and thymic activity predict lymphocyte recovery and outcomes after matched-sibling hematopoietic stem cell transplant. Haematologica. 2013 Mar; 98(3):346-52. < PMID: 23065508 >
- Atarashi K, Tanoue T, Oshima K, Suda W, Nagano Y, Nishikawa H, Fukuda S, Saito T, Narushima S, Hase K, Kim S, Fritz JV, Wilmes P, Ueha S, Matsushima K, Ohno H, Olle B, Sakaguchi S, Taniguchi T, Morita H, Hattori M, Honda K: Treg induction by a rationally selected mixture of Clostridiastrains from the human microbiota. Nature. 2013 Aug 8; 500(7461):232-6. < PMID: 23842501 >
- Blankenhaus B, Reitz M, Brenz Y, Eschbach ML, Hartmann W, Haben I, Sparwasser T, Huehn J, Kühl A, Feyerabend TB, Rodewald HR, Breloer M: Foxp3+ regulatory T cells delay expulsion of intestinal nematodes by suppression of IL-9-driven mast cell activation in BALB/c but not in C57BL/6 mice. PLoS Pathog. 2014 Feb 6; 10(2):e1003913. < PMID: 24516385 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com