

TECHNICAL DATA SHEET

CyFlow™ IgE PE-Cy5 Anti-Hu; Clone BE5

REF AM466973

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	IgE
Alternative Names	—
Clone	BE5
Clonality	monoclonal
Format	PE-Cy5
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [0.1 mg/ml]
Immunogen	Purified human IgE

Specificity

The mouse monoclonal antibody BE5 recognizes human immunoglobulin E (IgE); it recognizes an epitope different from the ones recognized by 4G7 and 4H10 antibodies to IgE.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Working concentrations should be determined by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH $\approx$ 7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

Immunoglobulin E (IgE) is a 180 kDa soluble protein serving as an antigen-specific unit of mast cell effector mechanisms. IgE has the lowest serum concentration of all immunoglobulins (approximately 0.5 mg/l) in healthy individuals, but upon allergen challenge its concentration in blood increases dramatically. Although biological survival of free IgE is very short ($T_{1/2}$ = 2 days), it is stabilized after binding to its high affinity receptor. Unlike IgM- IgG- and IgA-committed B cells, IgE-switched B cells do not undergo clonal expansion.

References

- Franklin EC: Structure and function of immunoglobulins. Acta Endocrinol Suppl (Copenh). 1975; 194:77-95. < PMID: 47690 >
- Fuller JM, Keyser JW: Serum immunoglobulins after surgical operation. Clin Chem. 1975 May; 21(6):667-71. < PMID: 1122610 >
- Balogh Z, Merétey K, Falus A, Bozsóky S: Serological abnormalities in juvenile chronic arthritis: a review of 46 cases. Ann Rheum Dis. 1980 Apr; 39(2):129-34. < PMID: 6966908 >
- Brinkmann V, Heusser CH: T cell-dependent differentiation of human B cells into IgM, IgG, IgA, or IgE plasma cells: high rate of antibody production by IgE plasma cells, but limited clonal expansion of IgE precursors. Cell Immunol. 1993 Dec; 152(2):323-32. < PMID: 8258141 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Gould HJ, Beavil RL, Vercelli D: IgE isotype determination: epsilon-germline gene transcription, DNA recombination and B-cell differentiation. Br Med Bull. 2000; 56(4):908-24. < PMID: 11359628 >
- Kaulfürst-Soboll H, Mertens M, Brehler R, von Schaewen A: Reduction of cross-reactive carbohydrate determinants in plant foodstuff: elucidation of clinical relevance and implications for allergy diagnosis. PLoS One. 2011 Mar 14; 6(3):e17800. < PMID: 21423762 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Cy and CyDye are trademarks of GE Healthcare.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com