

TECHNICAL DATA SHEET

CyFlow™ IgG (Fab) PE Anti-Hu; Clone 4A11

REF CZ138822

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	IgG (Fab)
Alternative Names	—
Clone	4A11
Clonality	monoclonal
Format	PE
Host / Isotype	Mouse / IgG2b
Species Reactivity	Human
Negative Species Reactivity	—
Quantity [Concentration]	0.1 mg [0.1 mg/ml]
Immunogen	Purified human IgG

Specificity

The mouse monoclonal antibody 4A11 recognizes Fab fragment of human immunoglobulin G (IgG).

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis. Suggested working usage is 6 µg/ml. Indicated dilution is recommended starting point for use of this product, but working concentrations should be validated by the investigator.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the total plasma immunoglobulins. IgG has the highest stability of blood immunoglobulins (T_{1/2} = 21 days) and is able of placental transfer. IgG is secreted by plasma cells at a comparably high rate as other immunoglobulins.

References

- Franklin EC: Structure and function of immunoglobulins. Acta Endocrinol Suppl (Copenh). 1975; 194:77-95. < PMID: 47690 >
- Fuller JM, Keyser JW: Serum immunoglobulins after surgical operation. Clin Chem. 1975 May; 21(6):667-71. < PMID: 1122610 >
- Balogh Z, Merétey K, Falus A, Bozsóky S: Serological abnormalities in juvenile chronic arthritis: a review of 46 cases. Ann Rheum Dis. 1980 Apr; 39(2):129-34. < PMID: 6966908 >
- Brinkmann V, Heusser CH: T cell-dependent differentiation of human B cells into IgM, IgG, IgA, or IgE plasma cells: high rate of antibody production by IgE plasma cells, but limited clonal expansion of IgE precursors. Cell Immunol. 1993 Dec; 152(2):323-32. < PMID: 8258141 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Wang H, Lee HK, Bukowski JF, Li H, Mariuzza RA, Chen ZW, Nam KH, Morita CT: Conservation of nonpeptide antigen recognition by rhesus monkey V gamma 2V delta 2 T cells.. J Immunol. 2003 Apr 1; 170(7):3696-706. < PMID: 12646635 >
- Fujii Y, Kaneko S, Nzou SM, Mwau M, Njenga SM, Tanigawa C, Kimotho J, Mwangi AW, Kiche I, Matsumoto S, Niki M, Osada-Oka M, Ichinose Y, Inoue M, Itoh M, Tachibana H, Ishii K, Tsuboi T, Yoshida LM, Mondal D, Haque R, Hamano S, Changoma M, Hoshi T, Kamo K, Karama M, Miura M, Hirayama K: Serological Surveillance Development for Tropical Infectious Diseases Using Simultaneous Microsphere-Based Multiplex Assays and Finite Mixture Models. PLoS Negl Trop Dis. 2014 Jul 31; 8(7):e3040. < PMID: 25078404 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com