

TECHNICAL DATA SHEET

CyFlow™ IgG1 Alexa Fluor™ 700 Mouse Isotype Control

REF AS153909

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	IgG1 Isotype Control
Alternative Names	—
Clone	MOPC-21
Clonality	monoclonal
Format	Alexa Fluor™ 700
Host / Isotype	Mouse / IgG1
Species Reactivity	n/a
Negative Species Reactivity	Human Rat
Quantity [Concentration]	0.1 mg [1 mg/ml]
Immunogen	< no data >

Specificity

This mouse IgG1 κ monoclonal antibody (clone MOPC-21) has unknown specificity and was chosen as an isotype control after screening on variety of resting, activated, live and fixed rat and human tissues.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed as an isotype control for Flow Cytometry analysis. To establish the amount of non-specific antibody binding, match the concentration of the correct isotype to the recommended working concentration of the antigen-specific antibody. If the background signal of the isotype control is too high (usually when working antibody concentrations are above 10 µg/ml of incubation mixture), change the experimental conditions to reduce the background.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

The specificity of staining by monoclonal antibodies to target antigens should be verified by establishing the amount of non-specific antibody binding. Especially at higher concentration (more than 15 µg/ml) the antibody staining usually has consignable background. To this end a non-reactive immunoglobulin of the same isotype is included as a negative control for each specific monoclonal antibody used in a particular immunoassay. The monoclonal antibody MOPC-21, generated against an undefined antigen, does not react specifically with rat and human samples, and hence all the background that could be observed when working with this antibody would be a result of general nonspecific interactions between an mouse IgG1 molecule and the respective sample under the particular conditions. This shall help the customer to set up the experimental conditions so that the nonspecific binding of any antibody is abolished.

References

- Wiendl H, Mitsdoerffer M, Schneider D, Melms A, Lochmuller H, Hohlfeld R, Weller M: Muscle fibres and cultured muscle cells express the B7.1/2-related inducible co-stimulatory molecule, ICOSL: implications for the pathogenesis of inflammatory myopathies. Brain. 2003 May; 126(5):1026-35. < PMID: 12690043 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Bryceson YT, March ME, Barber DF, Ljunggren HG, Long EO: Cytolytic granule polarization and degranulation controlled by different receptors in resting NK cells. J Exp Med. 2005 Oct 3; 202(7):1001-12. < PMID: 16203869 >
- Carlsten M, Björkström NK, Norell H, Bryceson Y, van Hall T, Baumann BC, Hanson M, Schedvins K, Kiessling R, Ljunggren HG, Malmberg KJ: DNAX accessory molecule-1 mediated recognition of freshly isolated ovarian carcinoma by resting natural killer cells. Cancer Res. 2007 Feb 1; 67(3):1317-25. < PMID: 17283169 >
- Yates J, Rovis F, Mitchell P, Afzali B, Tsang JY, Garin M, Lechler RI, Lombardi G, Garden OA: The maintenance of human CD4+ CD25+ regulatory T cell function: IL-2, IL-4, IL-7 and IL-15 preserve optimal suppressive potency in vitro. Int Immunol. 2007 Jun; 19(6):785-99. < PMID: 17545278 >
- Rebetz J, Tian D, Persson A, Widegren B, Salford LG, Englund E, Gisselsson D, Fan X: Glial progenitor-like phenotype in low-grade glioma and enhanced CD133-expression and neuronal lineage differentiation potential in high-grade glioma. PLoS One. 2008 Apr 9; 3(4):e1936. < PMID: 18398462 >
- Smed-Sörensen A, Moll M, Cheng TY, Loré K, Norlin AC, Perbeck L, Moody DB, Spetz AL, Sandberg JK: IgG regulates the CD1 expression profile and lipid antigen-presenting function in human dendritic cells via FcγRIIIa. Blood. 2008 May 15; 111(10):5037-46. < PMID: 18337560 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com