

TECHNICAL DATA SHEET

CyFlow™ Kappa-LC APC-Cy7 Anti-Hu; Clone TB28-2

REF BZ136230

For Research Use Only.

Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	κ light chains
Alternative Names	—
Clone	TB28-2
Clonality	monoclonal
Format	APC-Cy7
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	—
Quantity	100 tests
Immunogen	Human IgG-κ myeloma protein

Specificity

The mouse monoclonal antibody TB28-2 recognizes human Ig κ light chains of both secreted and cell surface immunoglobulin. It detects also free κ light chains.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

Immunoglobulin classes share the same basic four polypeptide chain structure of two heavy chains (five heavy chains types) and two light chains (κ and λ; both having a molecular weight of 22.5kDa). κ and λ consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of κ to λ is 70:30.

References

- Kiyotaki M, Cooper MD, Bertoli LF, Kearney JF, Kubagawa H: Monoclonal anti-Id antibodies react with varying proportions of human B lineage cells. J Immunol. 1987 Jun 15; 138(12):4150-8. < PMID: 3495581 >
- Nakamura T, Kubagawa H, Cooper MD: Heterogeneity of immunoglobulin-associated molecules on human B cells identified by monoclonal antibodies. Proc Natl Acad Sci USA. 1992 Sep 15; 89(18):8522-6. < PMID: 1382292 >
- Karandikar NJ, Aquino DB, McKenna RW, Kroft SH: Transient myeloproliferative disorder and acute myeloid leukemia in Down syndrome: An immunophenotypic analysis. Am J Clin Pathol. 2001 Aug; 116(2):204-10. < PMID: 11488066 >
- Böttcher S, Ritgen M, Buske S, Gesk S, Klapper W, Hoster E, Hiddemann W, Unterhalt M, Dreyling M, Siebert R, Kneba M, Pott C: EU MCL MRD Group: Minimal residual disease detection in mantle cell lymphoma: methods and significance of four-color flow cytometry compared to consensus

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

IGH-polymerase chain reaction at initial staging and for follow-up examinations. Haematologica. 2008 Apr; 93(4):551-9. < PMID: 18379010 >

- Jourdan M, Caraux A, De Vos J, Fiol G, Larroque M, Cognot C, Bret C, Duperray C, Hose D, Klein B: An in vitro model of differentiation of memory B cells into plasmablasts and plasma cells including detailed phenotypic and molecular characterization. Blood. 2009 Dec 10; 114(25):5173-81. < PMID: 19846886 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Cy and CyDye are trademarks of GE Healthcare.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com