

TECHNICAL DATA SHEET

CyFlow™ Kappa-LC PE Anti-Hu; Clone A8B5

REF CV648717

For Research Use Only.
Not for use in diagnostic or therapeutic procedures.

Specifications

Antigen	κ light chains
Alternative Names	—
Clone	A8B5
Clonality	monoclonal
Format	PE
Host / Isotype	Mouse / IgG1
Species Reactivity	Human
Negative Species Reactivity	Sheep Goat Rabbit Guinea pig Hamster
Quantity	100 tests
Immunogen	< no data >

Specificity

The mouse monoclonal antibody A8B5 recognizes κ light chains (22.5 kDa) of immunoglobulins.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

Application

The reagent is designed for Flow Cytometry analysis of human blood cells. Recommended usage is 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.

Other usages may be determined from the scientific literature.

Storage Buffer

The reagent is provided in stabilizing phosphate buffered saline (PBS) solution, pH ≈7.4, containing 0.1% (w/v) sodium azide.

Storage and Stability

Storage	Avoid prolonged exposure to light. Store in the dark at 2-8°C. Do not freeze.
Stability	Do not use after expiration date stamped on vial label.

Background Information

Immunoglobulin classes share the same basic four polypeptide chain structure of two heavy chains (five heavy chains types) and two light chains (κ and λ; both having a molecular weight of 22.5kDa). κ and λ consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of κ to λ is 70:30.

References

- Franklin EC: Structure and function of immunoglobulins. Acta Endocrinol Suppl (Copenh). 1975; 194:77-95. < PMID: 47690 >
- Diaz de Domingo NB, Lardo MM, Romero Artaza J, Merelli A, Carbia CD, Sánchez Avalos JC: Aplicacion de tecnicas inmunocitoquimicas para el diagnostico de neoplasias hematologicas. Medicina (B Aires). 1999; 59(1):11-6. < PMID: 10349112 >
- Konig A, Wennemuth G, Soyer HP, Hoffmann R, Happel R, Krause W: Vulvar amyloidosis mimicking giant condylomata acuminata in a patient with multiple myeloma. Eur J Dermatol. 1999 Jan-Feb; 9(1):29-31. < PMID: 9920983 >
- Scheeren FA, Diehl SA, Smit LA, Beaumont T, Naspetti M, Bende RJ, Blom B, Karube K, Ohshima K, van Noesel CJ, Spits H: IL-21 is expressed in Hodgkin lymphoma and activates STAT5: evidence that activated STAT5 is required for Hodgkin lymphomagenesis. Blood. 2008 May 1; 111(9):4706-15. < PMID: 18296629 >

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com

- Di Bartolo DL, Hyjek E, Keller S, Guasparri I, Deng H, Sun R, Chadburn A, Knowles DM, Cesarman E: Role of defective Oct-2 and OCA-B expression in immunoglobulin production and Kaposi's sarcoma-associated herpesvirus lytic reactivation in primary effusion lymphoma. J Virol. 2009 May; 83(9):4308-15. < PMID: 19224997 >
- Avilés-Salas A, Chanona-Vilchis JG, Candelaria-Hernández MG, Peña-Torres Mde L, Cuéllar-Mendoza ME: Presentación inicial de un linfoma plasmablastico como tumor ovárico. Ginecol Obstet Mex. 2012 Jun; 80(6):430-4. < PMID: 22826973 >

The Safety Data Sheet for this product is available at www.sysmex-partec.com/services.

Contact Information:

Sysmex Partec GmbH • Am Flugplatz 13 • 02828 Görlitz • Germany
Tel +49 3581 8746 0 • Fax +49 3581 8746 70 • E-mail: info@sysmex-partec.com